



PROTOCOL

HVTN 805/HPTN 093

**Antiretroviral analytical treatment interruption (ATI) to
assess immunologic and virologic responses in
participants who initiated ART in early HIV infection
after having received VRC01 or placebo in HVTN
703/HPTN 081**

DAIDS DOCUMENT ID 38691

Non-IND Protocol

CLINICAL TRIAL SPONSORED BY

Division of AIDS (DAIDS)
National Institute of Allergy and Infectious Diseases (NIAID)
National Institutes of Health (NIH)
Department of Health and Human Services (DHHS)
Bethesda, Maryland, USA

April 27, 2020
HVTN 805/HPTN 093
Version 1.0

Contents

1	Protocol summary	4
1.1	Précis	6
1.2	Schema	7
1.3	Protocol team.....	8
2	Background and rationale	10
2.1	Antiretroviral therapy	10
2.2	Early ART initiation is associated with delayed time to rebound and post-treatment control	10
2.3	Broadly neutralizing antibodies (bnAbs)	15
2.4	Antiretroviral analytical treatment interruption (ATI)	26
2.5	An AMP ATI.....	33
2.6	Ethical considerations	34
2.7	Feasibility	35
2.8	Conclusion.....	37
3	Study design.....	38
3.1	Hypotheses	38
3.2	Objectives and endpoints	38
3.3	ATI	41
4	Statistical considerations.....	45
4.1	Estimated accrual	45
4.2	Statistical analyses.....	46
5	Study population	51
5.1	Inclusion criteria.....	51
5.2	Exclusion criteria.....	53
5.3	Criteria for participant early termination.....	55
5.4	Study termination	56
6	Study procedures.....	57
6.1	Schedule 1: Monitoring ATI	57
6.2	Schedule 2: Monitoring ATI with viremia	62
6.3	Schedule 3: Follow-up on ART	63
6.4	Interim visits.....	64
6.5	Early termination visit.....	65
6.6	Contraception status	65
6.7	Specific clinical management considerations	66
7	Study products	67
8	Laboratory	68
8.1	CRS laboratory procedures	68
8.2	Assay timepoints	68
8.3	Endpoint assays: cellular	68
8.4	Phenotyping of cell populations	69
8.5	Endpoint assays: humoral.....	69
8.6	HIV viral reservoir	69
8.7	Host genomics	70

8.8	HIV antiretroviral resistance testing.....	70
8.9	ARV detection.....	70
8.10	Biohazard containment.....	70
9	Research use of stored human samples and/or data.....	71
9.1	Specimen storage and other use of specimens	71
9.2	Exploratory studies.....	71
10	Potential risks and benefits	72
10.1	Potential risks	72
10.2	Potential benefits.....	72
11	Monitoring of safety	73
11.1	Safety monitoring and oversight	73
11.2	Safety reporting	75
12	Ethics/protection of human subjects	77
12.1	Informed consent process.....	77
12.2	Participant confidentiality	78
13	Protocol conduct	79
13.1	Emergency communication with study participants	80
14	Acronyms and abbreviations.....	81
15	Protocol version history	83
16	Document references (other than literature citations).....	84
17	References.....	86
Appendix A	Sample informed consent form	97
Appendix B	Approved birth control methods (for sample informed consent form)	120
Appendix C	Sample consent form for use of samples and information in other studies.....	121
Appendix D	Tables of procedures for sample informed consent form.....	126
Appendix E	Laboratory procedures—Schedule 1: Monitoring ATI.....	131
Appendix F	Laboratory procedures—Schedule 2: Monitoring ATI with viremia.....	135
Appendix G	Laboratory procedures—Schedule 3: Follow-up on ART	139
Appendix H	Procedures at CRS—Schedule 1: Monitoring ATI.....	141
Appendix I	Procedures at CRS—Schedule 2: Monitoring ATI with viremia.....	145
Appendix J	Procedures at CRS—Schedule 3: Follow-up on ART	147
Appendix K	Visit Windows.....	148
Appendix L	Protocol Signature Page	153

1 Protocol summary

Full title:	Antiretroviral analytical treatment interruption (ATI) to assess immunologic and virologic responses in participants who initiated ART in early HIV infection after having received VRC01 or placebo in HVTN 703/HPTN 081
Short title:	HVTN 805/HPTN 093
Sponsor:	NIAID Division of AIDS
Conducted by:	HIV Vaccine Trials Network (HVTN) and HIV Prevention Trials Network (HPTN)
Protocol chairs:	Shelly Karuna, MD, MPH; Katharine Bar, MD; Simba Takuva, MBChB, MSc
Sample size:	16 - 61
Study population:	HVTN 703/HPTN 081 participants living with HIV who met criteria for transition to Schedule 2 or Schedule 3 in that trial
Study design:	An exploratory study of participants living with HIV undergoing an analytical treatment interruption after early initiation of antiretroviral therapy (ART) following HIV acquisition in HVTN 703/HPTN 081, where they received VRC01 or placebo infusions
Study duration:	Study duration is potentially indefinite for a participant maintaining extended viral control during ATI. Study duration for most participants is expected to be 13-18 months. The maximum anticipated duration for any participant is expected to be approximately 2½ to 3 years.
Study products:	None. Drugs for ART and for pre-exposure prophylaxis (PrEP) will not be provided by the study or paid for using sponsor funds. Access to external funding sources for PrEP and ART provision is available and is detailed in the HVTN 805/HPTN 093 Study Specific Procedures (SSP).
Primary hypotheses:	Individuals who initiated ART early in HIV infection in HVTN 703/HPTN 081 will suppress plasma viremia and maintain CD4+ T-cell counts longer during ATI than historical cohorts (ie, SPARTAC African and non-African cohorts described in Gossez et al (1) comprising individuals who initiated ART early in HIV infection without other interventions).

Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will suppress plasma viremia and maintain CD4+ T-cell counts longer during ATI than those who initiated ART early in HIV infection and received placebo within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081.

ATI will be safe and well-tolerated in individuals who acquired HIV during their participation in HVTN 703/HPTN 081.

Primary objectives: To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection, on the time to meeting ART re-initiation criteria in participants undergoing ATI

To evaluate the safety of ATI among HVTN 805/HPTN 093 participants

Secondary hypotheses: Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will have enhanced cellular and humoral responses compared with those who initiated ART early in HIV infection and received placebo within 8 weeks of acquiring HIV in HVTN 703/HPTN 081.

Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will have more limited viral reservoirs, before and after ATI, than those who initiated ART early in HIV infection and received placebo within 8 weeks of acquiring HIV in HVTN 703/HPTN 081.

Secondary objectives: To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection, on the development of anti-HIV immune responses and on the potential association of those immune responses with time to meeting criteria for ART re-initiation in participants undergoing ATI

To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV

acquisition period and/or during early infection, on viral load in participants undergoing ATI

To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection, on HIV reservoir size before and after ATI, and whether HIV reservoir measurements are associated with time to meeting criteria for ART re-initiation in participants undergoing ATI


1.1 Précis

While plasma viremia rebounds rapidly in most chronically HIV-infected individuals upon cessation of therapy, early initiation of ART after HIV diagnosis has been associated with later ART-free virologic control in individuals known as “post-treatment controllers” (PTCs). Similarly, preclinical (eg, murine, nonhuman primate [NHP]) and clinical evidence suggest that the presence of broadly neutralizing anti-HIV-1 antibodies (bnAbs), even at a concentration, or with a neutralization sensitivity profile, insufficient to prevent HIV acquisition, may nevertheless modulate the autologous response to HIV infection in ways that help set the stage for later durable virologic control. There is also evidence suggesting that early ART initiation and the presence of bnAbs at the time of or soon after HIV infection may be capable of limiting the establishment of and decreasing the size of viral reservoirs. Whether and how these effects might act together to support durable viral control, and what biomarkers may predict such control, is unknown.

The Antibody Mediated Prevention (AMP) studies HVTN 704/HPTN 085 and HVTN 703/HPTN 081, phase 2b studies assessing the ability of the CD4 binding site bnAb VRC01 to prevent HIV-1 infection, provide a unique opportunity to assess the impact of these hypothesized effects. The AMP studies currently have the only cohort of HIV-1–infected individuals who were diagnosed soon after acquisition, who initiated ART relatively soon after diagnosis, and a subset of whom had bnAb present at or near the time of infection. A pause in antiretroviral treatment—that is, an ATI—is the only way to determine whether early ART initiation with and without VRC01 circulating around the time of HIV acquisition, results in blunted, delayed, or absent viral rebound following discontinuation of ART, or subsequent immune control of any rebound that may be observed. If such an impact is observed, ATI is also the best means of assessing mechanisms and identifying predictive biomarkers of that impact through assessment of innate, cellular, and humoral markers along with characterization of the viral reservoir. Extensive virologic, immunologic, and safety monitoring will be conducted, and the trial has been formulated for consistency with recent consensus guidelines on the design and conduct of ATI studies (2, 3), particularly with respect to risk mitigation.

The study schema is shown below.

1.2 Schema



			SCHEDULE 1			SCHEDULE 2				SCHEDULE 3		
	SCREEN	PRE-DISCONTINUE ART	ATI WEEKS 0-8	ATI WEEKS 10-24	ATI WEEKS 28-52 ¹	ATI + Viremia WEEKS 0-8	ATI + Viremia WEEKS 10-36	ATI + Viremia WEEKS 40-52 ²	PRE-REINITIATE ART	FOLLOW UP ON ART WEEKS 0-12	FOLLOW UP ON ART WEEKS 12-28	FOLLOW UP ON ART WEEKS 40-52
Plasma HIV RNA	✓	✓	WEEKLY	Q2 WEEKS	Q4 WEEKS	WEEKLY	Q2 WEEKS ³	Q4 WEEKS	✓	Q2 WEEKS	Q4 WEEKS	Q12 WEEKS
CD4+ & CD8+ T cell counts	✓	✓	Q2 WEEKS	Q4 WEEKS	Q8 WEEKS	Q2 WEEKS	Q4 WEEKS ⁴	Q8 WEEKS	✓	Q4 WEEKS	Q8 WEEKS	Q12 WEEKS
Hematology & Chemistries	✓	✓	Q4 WEEKS		Q8 WEEKS	Q4 WEEKS		Q8 WEEKS	—	Q4 WEEKS	Q12 WEEKS	Q12 WEEKS

¹ QUARTERLY FOLLOW-UP VISITS MAY CONTINUE BEYOND WEEK 52 FOR PARTICIPANTS WHO DO NOT MEET CRITERIA FOR TRANSITION TO SCHEDULE 2.

² QUARTERLY FOLLOW-UP VISITS MAY CONTINUE BEYOND WEEK 52 FOR PARTICIPANTS WHO DO NOT MEET CRITERIA FOR ART RE-INITIATION.

³ OR WEEKLY FOR WEEKS 10-24, IF VL ≥ 200 copies/mL

⁴ OR Q2 WEEKS FOR WEEKS 10-24 IF VL ≥ 200 copies/mL

1.3 Protocol team

Protocol leadership

<i>Co-chairs</i>	Shelly Karuna HVTN Core, Fred Hutch 206-667-4355 skaruna@fredhutch.org	<i>Medical officers</i>	Randall Tressler DAIDS Therapeutics Research Program 240-627-3072 randall.tressler@nih.gov
	Katharine Bar Univ. of Pennsylvania 215-573-8497 bark@pennmedicine.upenn.edu		David Burns DAIDS Prevention Sciences Program (PSP), NIAID 301-435-8896 burnsda@niaid.nih.gov
	Simba Takuva HVTN Core, Fred Hutch 011-989-9754 stakuva@hcrisa.org.za		Margarita Gomez Lorenzo DAIDS Vaccine Research Program (VRP), NIAID 240-627-3510 gomezmm@niaid.nih.gov
<i>Protocol Team leaders</i>	Shelly Karuna HVTN Core, Fred Hutch 206-667-4355 skaruna@fredhutch.org	<i>Laboratory leads</i>	John Hural HVTN Laboratory Program, Fred Hutch 206-667-1683 jhural@fredhutch.org
	Phil Andrew HPTN LOC, FHI 360 919.544.7040 x11213 pandrew@fhi360.org		Estelle Piwowa-Manning HPTN Laboratory Center Johns Hopkins University 410-614-6736 epiwowa@jhmi.edu
<i>Statistician</i>	Allan DeCamp SCHARP, Fred Hutch 206-667-7892 adecamp@scharp.org		

Other contributors

<i>Core medical monitor</i>	Shelly Karuna HVTN Core, Fred Hutch	<i>HVTN Regional medical liaison, Africa</i>	Simba Takuva HVTN Core, Fred Hutch
<i>Consulting investigators</i>	Tae-Wook Chun NIAID Laboratory of Immunoregulation	<i>HVTN Clinical safety specialist</i>	Maija Anderson HVTN Core, Fred Hutch
	Lucio Gama NIAID Vaccine Research Center	<i>HVTN Social/behavioral scientist</i>	Michele Andrasik HVTN Core, Fred Hutch
	Catherine Orrell Desmond Tutu HIV Centre, Cape Town, South Africa	<i>HVTN Clinical trials manager</i>	Carissa Karg HVTN Core, Fred Hutch
	Nyaradzo Mgodi UZ-UCSF Collaborative Research Programme Harare, Zimbabwe	<i>HPTN Clinical research manager</i>	Phil Andrew HPTN LOC, FHI 360
<i>HVTN Laboratory representative</i>	Lisa Sanders HVTN Laboratory Program, Fred Hutch	<i>Statistical research associates</i>	Erika Rudnicki SCHARP, Fred Hutch

<i>HPTN Laboratory representative</i>	Vanessa Cummings HPTN Laboratory Center Johns Hopkins University		Chenchen Yu SCHARP, Fred Hutch
<i>Regulatory affairs associate</i>	Megan Brandon HVTN Core, Fred Hutch	<i>Clinical data manager</i>	Alison Ayres SCHARP, Fred Hutch
<i>Clinic coordinators</i>	Nitesha Jeenarain South Africa MRC Chatsworth CRS Durban, South Africa TBD	<i>SDMC Associate director of lab science</i>	April Randhawa SCHARP, Fred Hutch
		<i>Protocol development manager</i>	Carter Bentley HVTN Core, Fred Hutch
<i>Community Advisory Board (CAB) members</i>	Angy Peter Groote Schuur CAB Cape Town, South Africa Maximina Jokonya University of Zimbabwe College of Health Sciences Clinical Trials Research Center (UZCHS-CTRC) CAB Harare, Zimbabwe	<i>HVTN Community engagement unit representatives</i>	Gail Broder HVTN Core, Fred Hutch Kagisho Baepanye HVTN Core Johannesburg, South Africa
<i>Prevention research specialist</i>	Michelle Robinson HPTN LOC, FHI 360	<i>HPTN Community program associate</i>	Jonathan Lucas HPTN LOC, FHI 360
<i>Ethics representative</i>	Ames Dhai University of the Witwatersrand, Johannesburg, South Africa	<i>Community educator/recruiters</i>	Charles Chasakara University of Zimbabwe College of Health Sciences Clinical Trials Research Center (UZCHS-CTRC) Harare, Zimbabwe Ivy Kaunda CAPRISA-eThekweni Durban, South Africa

2 Background and rationale

2.1 Antiretroviral therapy

Antiretroviral therapy (ART) can now achieve prolonged suppression of plasma viremia in most HIV-infected individuals. Consequently, ART has dramatically improved the mortality, morbidity, quality of life, and overall clinical course of infected individuals. However, clinically successful suppression of HIV comes at a cost; long-term toxicity, drug-drug interactions, stigma, drug resistance, adherence challenges, and pill fatigue necessitate a continued search for effective alternatives for achieving durable control of HIV replication in infected individuals. Furthermore, ART alone has been unable to completely eradicate HIV; plasma viremia rebounds rapidly with reactivation of persistent viral reservoirs in almost all chronically HIV-infected individuals upon cessation of therapy (4-7). Persistence of HIV reservoirs, even with successful suppression of plasma viremia by ART, is a major obstacle to HIV eradication.

Thus, one of the highest priorities for the HIV field is the search for therapeutic strategies that can eliminate or control HIV replication and reservoirs in the absence of ART. Early initiation of ART after HIV diagnosis has been associated with later ART-free virologic control in individuals known as “post-treatment controllers” (PTCs). Another potential strategy is to harness the immune system of people living with HIV to increase their ability to control HIV off ART; broadly neutralizing monoclonal antibodies (bnAbs) against HIV, present at the time of HIV acquisition, may help increase the immune system’s capacity to contribute to ART-free control.

2.2 Early ART initiation is associated with delayed time to rebound and post-treatment control

There is a substantial body of evidence suggesting early initiation of ART is associated with both delayed time to rebound after ATI and durable post-treatment control. PTCs are generally considered to be individuals who, after maintaining some period of viral suppression on ART, stop ART and maintain virologic control for a prolonged period of time, generally more than 6 to 12 months. Investigators are also interested in shorter durations of virus suppression, which are captured by studies comparing time to rebound in the immediate weeks to months after treatment interruption. It is unclear to what degree durable post-treatment control and more transient periods of delayed time to rebound are related mechanistically, but both phenomena are enriched in people living with HIV with early ART initiation.

2.2.1 Delayed time to rebound

Multiple clinical trials have shown that individuals with ART initiation during early infection (generally after antibody seroconversion but within ~6 months of infection, or Fiebig stages III-V), and to a lesser degree, during acute infection (generally pre-antibody seroconversion, or Fiebig stages I-II) experience delayed viral rebound compared to individuals starting antiretrovirals in chronic infection. In a metanalysis of 6 AIDS Clinical Trials Group (ACTG) trials compiling data from individuals who underwent ATI without intervention, Jonathan Li and colleagues (8) found that participants starting ART during acute infection (n = 32) and early infection (n = 48) maintained virus suppression at 12 weeks post-ATI at increased rates compared to those with ART initiation during chronic infection (n = 155) (9% and 15% vs 3%, respectively, $p = 0.01$, Fisher's exact test) (see [Figure 2-1](#)). As possible mechanisms for the longer time to rebound, the study explored issues of size and transcriptional activity of the participants' viral reservoirs. In subset analyses, participants with acute and early ART initiation (n = 20) were found to have less transcriptionally active virus reservoirs than participants with chronic ART initiation (n = 104) (cell-associated RNA (caRNA) levels of 92 vs 156 copies/ 10^6 CD4 T cells, $p < 0.01$, exact Wilcoxon) and a statistically non-significant trend toward smaller reservoirs, as measured by total HIV DNA. Notably, these 6 ACTG trials enrolled predominantly white men from the United States infected with clade B viruses (91% men, 77% Caucasian).




Figure 2-1 Virologic suppression after treatment interruption stratified by timing of ART initiation. (b) Cumulative percentage of participants who remained suppressed stratified by treatment initiated during acute, early, or chronic HIV infection at the 200 HIV ribonucleic acid (RNA) copies/ml and (c) 1000 HIV RNA copies/ml viral rebound thresholds. From (8), Figure 1.

Several other studies have corroborated the effect of early ART initiation on delayed time to rebound. For example, Steingrover et al (9) described results from trials of ATI in individuals initiating ART during primary HIV infection (PHI) (n = 24) vs chronic HIV infection (CHI) (n = 48). Participants with early ART initiation rebounded later than those with chronic ART initiation (median 8 weeks vs 4 weeks, with viral rebound thresholds of > 50, > 500, and > 5000 copies/ml)

(Figure 2-2). In the early ART group, 4 participants (17%) maintained virus suppression < 50 copies/mL at 48 weeks, while no chronic ART initiators maintained suppression. This European trial enrolled largely white men (87% men, 76% Western origin) and results are further limited by frequent use of non-nucleoside reverse transcriptase inhibitor (NNRTI) antiretrovirals with long half-lives and relatively infrequent sampling.




Figure 2-2 Longitudinal plots of plasma HIV-1 RNA after treatment interruption. (a) The mean plasma HIV-1 RNA load plotted against the time in weeks after interruption of HAART (stop) for CHI (squares) and PHI (triangles) groups. Error bars represent 95% confidence intervals. Kaplan-Meier plots showing the different time to plasma viral rebound for CHI and PHI groups. Rebound of plasma HIV-1 viral load was defined as a single measurement of plasma HIV-1 RNA above (a) 5000 copies/ml, (b) 500 copies/ml or (c) 50 copies/ml. Differences between the groups were tested using the log rank test. CHI, chronic HIV infection; HAART, highly active antiretroviral therapy; PHI, primary HIV infection. From (9), Figure 1.

As noted above, the literature describing viral kinetics after treatment interruption is limited in women and non-Westerners. A recent publication describing treatment interruption in the PROMISE trial provides one of the only analyses of young, international, post-partum women. Le et al monitored viral kinetics and safety after a treatment interruption in 1,076 virally suppressed, post-partum women (82% Black/African, 7% White, 7% Asian/Pacific Islander) (10). The vast majority of PROMISE participants rebounded early (median 2 weeks post-ATI); however, in comparison to the largely male, American cohort described above in the ACTG metanalysis (8), the PROMISE cohort had a significantly higher proportion of virally suppressed participants at 12 weeks post-ATI (25.4% vs

6.4%, $p < 0.0001$). A comparison between the PROMISE women and the American women in the ACTG study ($n = 22$) demonstrated a non-significant trend in the same direction (25.4% vs 13.6%, $p = 0.11$). Importantly, the ATI was well tolerated with very few adverse events. This study is partially limited by missing data points, use of longer-lived NNRTI antiretrovirals, and confounding immunologic effects of recent pregnancy. This cohort is also missing information on timing of ART initiation; however, it highlights the potential role of sex and epidemiology (including host genetic, environmental, or virus subtype differences) in virologic control post-ATI.

2.2.2 Post-treatment control

Evidence is similarly strong that early ART initiation is associated with the phenomenon of durable post-treatment control. While post-treatment control has been observed at different rates across cohorts, it is consistently more frequent in groups of early ART initiators (see [Figure 2-3](#); note that follow-up of participants in some of these cohorts continues, hence increases in duration of virologic control is expected over time).




Figure 2-3 Estimates of the frequency of post-treatment control and remission. Each data point represents an individual study, which has measured the frequency of post-treatment control or remission following treatment interruption among individuals who commenced ART during primary HIV infection (PHI). Included are analyses of the following studies: CASCADE (11, 12), ANRS Primo (13), VISCONTI (14, 15), SPARTAC (16, 17), CHAMP (18), ZHPI (19), SeaPIP (20) and NCT01859325 (21). Those which define post-treatment control (here considered to be > 24 months following treatment interruption) are colored green, those defining remission (> 4 months following treatment interruption) are shown in red. In some cases, a study explored both phenotypes; where possible these are disaggregated and listed twice, otherwise these are shown as remission. The duration of control is the time between

treatment interruption and viral rebound, or the end of follow-up, and is plotted at the median value amongst controllers. The size of the point corresponds to the median duration of ART prior to treatment interruption amongst controllers. Where a study did not report either of these values, the minimum value for inclusion is used instead. From (22) Figure 1.

The VISCONTI cohort is one the first and the best characterized groups of PTCs. This cohort arose from the French ANRS PRIMO study of early ART initiation, which identified 14 of 164 participants (9%) who maintained virus suppression after a median of 4.5 years of follow up (13). The 14 PTCs were subsequently found to have distinct characteristics from elite controllers (individuals who achieve virus control prior to ART initiation), including small but detectable latent reservoirs, few protective human leukocyte antigen (HLA) alleles, weak HIV-specific CD8⁺ T-cell responses, and more severe acute retroviral syndromes (ARS) (14, 18, 23-25).

The characteristics of the VISCONTI cohort and participants with delayed time to rebound give some hints into the potential mechanisms underlying virus suppression post-ATI and why it is more frequent with early ART. In general, early ART initiators have smaller, less diverse, and less transcriptionally active reservoirs. Further, their immune systems are believed to be better preserved and less exhausted. A recent sub-study of the SPARTAC trial has borne out some of these hypotheses.

The SPARTAC trial, which evaluated the impact of immediate vs delayed ART initiation after HIV diagnosis in both European and African participants, reported a 14% rate of post-treatment control at 1 year (16). Initial analyses suggested that smaller reservoirs were associated with PTC, as levels of total HIV-1 DNA prior to ART initiation and at ATI were associated with time to rebound. More recently, an in-depth analysis of the immunologic characteristics of the European (Subtype B virus-infected) SPARTAC PTCs has shown that measures of T-cell exhaustion (PD-1, Lag-3, and TIM-3) prior to ART initiation are strongly correlated with both levels of total HIV-1 DNA and time to rebound (26). Notably, no biomarkers at the time of ATI were identified that correlated with time to rebound. These results confirm the importance of smaller reservoirs and functional immune systems in post-treatment control, but do not elucidate a simple mechanism nor shed light on testing that can be done on ART-suppressed individuals to better predict their response to ATI.

Post-treatment control has continued to be reported in clinical trials and observational cohorts. Like the increased frequency of delayed rebound seen in African women in the PROMISE study, durable post-treatment control may be more common among African women. A recent SPARTAC sub-study revealed that the African SPARTAC cohort, all females infected with non-B (primarily C) subtypes, was more likely than the non-African SPARTAC cohort to maintain virologic control at study end (HR 3.9, 95% CI 1.75-8.81, $p < 0.001$) (1, 17). Further, 5 of 22 (23%) maintained VL <400 copies for the duration of follow up (median 4.48 years), while none of the non-African men maintained VL < 400 for

this duration. Of note, this frequency of post-treatment control is greater than any others included in the Martin et al analysis shown above in [Figure 2-3](#).

Gossez and colleagues reported a more in-depth analysis of the controllers, which is relevant to this proposal for several reasons (1). First, the cohort of African women studied in SPARTAC is similar to the potential participants HVTN 805/HPTN 093, in that they were young southern African women (91% from South Africa, median age of 26 years [interquartile range (IQR) 21- 31.8]), infected with predominantly non-clade B viruses (86.5% clade C), with early ART initiation (median 15.5 [IQR 13.2-17.7] weeks between seroconversion to ART initiation). ATI was safe and well-tolerated in SPARTAC and the prolonged ATI experienced by the 5 African women achieving post-treatment control was also safe and well tolerated, with CD4 counts remaining high (median 962 cells/ul [IQR 840-1028]) and reservoir measures remaining stable (total DNA, median 3.317 copies/million CD4 T cells [IQR 3.03-3.18]). Three of the 5 post-treatment controllers had episodes of transient viremia (ie, viral loads > 400 copies/mL, between 560-1690 copies/mL), supporting more permissive ART restart criteria. Finally, reservoir and immunologic characterizations were performed, but no predictor of delayed rebound or post-treatment control was identified, reinforcing the dearth of biomarkers and the current need for ATI to assess whether PTC will occur.

2.3 Broadly neutralizing antibodies (bnAbs)

Over the past decade, several bnAbs have been discovered and isolated and their HIV-1 target sites and mechanisms of neutralization have been elucidated (27-34). These efforts have informed the development of recombinant HIV vaccine immunogens designed to elicit broadly neutralizing antibodies (31, 35-39) and they have set the stage for an evaluation of these antibodies' potential as tools for HIV-1 prevention (ie, passive immunization or antibody-mediated prevention), treatment, and possibly cure (40-45).

Several early phase trials in HIV-uninfected and HIV-infected individuals demonstrated the safety, tolerability, antiviral activity, and PK of VRC01 (46-50) and in 2016 VRC01 advanced to the first bnAb prevention efficacy trials in HIV-uninfected individuals, the Antibody-Mediated Prevention (AMP) trials HVTN 704/HPTN 085 and HVTN 703/HPTN 081.

2.3.1 The HVTN 703/HPTN 081 AMP study

HVTN 703/HPTN 081 opened in May 2016 and enrolled 1924 women (persons assigned female sex at birth) at risk of HIV infection. Participants were randomized 1:1:1 in a double-blind fashion to receive 10 mg/kg VRC01, 30 mg/kg VRC01, or placebo. Participants receive study product via intravenous (IV) infusion every 8 weeks for a total of 10 infusions over 72 weeks and are then followed every 4 weeks to week 80, and every 8 weeks thereafter to week 104

(51) (see [Table 2-1](#)). In addition to assessing the prevention efficacy of VRC01 and providing further VRC01 safety and PK data, the AMP studies include assessment of functional activity (ie, fragment crystallizable (Fc) effector functions), sieve analyses, and correlates analyses. HVTN 703/HPTN 081 completed enrollment in October 2018.

Table 2-1 HVTN 703/HPTN 081 AMP schema

			Infusion schedule (Weeks)											
			[A = VRC01 infusion; C = Control infusion]											
	Treatment	N*	W0	W8	W16	W24	W32	W40	W48	W56	W64	W72	W80*	W104†
Group 1	VRC01 10 mg/kg	633	A	A	A	A	A	A	A	A	A	A		
Group 2	VRC01 30 mg/kg	633	A	A	A	A	A	A	A	A	A	A		
Group 3	Control	634	C	C	C	C	C	C	C	C	C	C		

* Due to the randomization scheme, the numbers of VRC01 and control recipients may differ slightly.

♦ Week 80 is the last study visit for the primary endpoint analysis of prevention efficacy.

† Week 104 is the last study visit for the co-primary endpoint analysis of safety and tolerability.

HIV diagnostic testing is conducted every 4 weeks and is accompanied by collection of serum and, at up to 4 timepoints, peripheral blood mononuclear cells (PBMCs) for storage. Participants who acquire HIV discontinue further study infusions and are monitored for 24 weeks with viral loads, CD4+ T-cell counts, and VRC01 levels, while providing serum, plasma and PBMCs for storage, functional humoral assays and other assessments. HIV-infected participants are linked to care for ART initiation and additional clinical follow-up.

Most HVTN 703/HPTN 081 participants initiate ART within 4 weeks of confirmation of HIV infection, a mean of 7 weeks after the first RNA-positive HIV diagnostic sample. The vast majority of participants exhibit HIV diagnostic patterns of RNA-positive, enzyme immunoassay (EIA)-positive, and Bio-Rad Geenius-negative, -indeterminate, or -positive with p31-negative at the time of their confirmatory HIV diagnostic sample, and initiate ART within 4 weeks of this time point.

2.3.2 bnAbs can mediate virologic control

The mission of the HVTN and of the HPTN is to prevent HIV, and the aim of the AMP studies is to evaluate the prevention efficacy of the VRC01 bnAb. However, for those participants who acquire HIV in spite of their participation in HVTN 703/HPTN 081, the potential impact of bnAbs beyond prevention efficacy is important to evaluate. Preclinical (eg, murine, NHP) and clinical (ie, human) evidence is mounting that anti-HIV bnAbs like VRC01 can mediate sustained virologic control of HIV.

For instance, in a preclinical experiment designed to assess the impact of early immunotherapy postacquisition, VRC01 was administered 10 days after simian-human immunodeficiency virus (SHIV)_{SF162P3} inoculation (ie, immediately prior to projected peak viremia) in 6 rhesus macaques, while another 6 macaques received a combination of VRC07-523LS + PGT121, and another 6 macaques received ART alone from day 10. Though SHIV_{SF162P3} is ~10-fold less sensitive to VRC01 than most isolates and both VRC07-523LS and PGT121 are more potent than VRC01, suppression of viral replication was comparable across all groups prior to the development of anti-monoclonal antibody (mAb) antibodies, a known limitation of the NHP model for bnAb administration (see [Figure 2-4](#)) (52).




Figure 2-4 Geometric mean plasma SHIV viral loads (pVL) for the untreated group and the 3 treatment groups (n = 6 per group). The error bars represent the 95% confidence intervals at each time point. From (52) Figure 2B.

In another recent study (53), macaques were infected with SHIV_{AD8-EO}. Three days after infection (prior to antibody seroconversion), the animals were randomized to receive either a 2-week course of 10-1074 and 3BNC117 or standard ART. At 15 weeks postinfection, the ART group discontinued ART. Six of the 13 animals treated with 10-1074 and 3BNC117 exhibited prolonged suppression of plasma viremia long after serum levels of infused antibodies became undetectable; in 5 of these controller macaques, this period of suppression followed an initial rebound of viremia with preservation of CD4⁺ T cells (see [Figure 2-5](#) and [Figure 2-6](#)). Although the 7 noncontroller bnAb-treated macaques did not fully suppress virus, 4 of those 7 maintained CD4⁺ T cell counts and low levels of viremia (105 – 385 copies/mL) without ART for 2-3 years postinfection. Thus, 10 of the 13 bnAb-treated macaques appeared to benefit from early immunotherapy. In contrast, none of the 3 ART-treated animals exhibited sustained suppression of plasma viremia following discontinuation of ART. No significant differences in HIV-specific CD8⁺ or CD4⁺ T-cell responses or anti-gp120 binding antibody levels were observed in controller and noncontroller bnAb-treated macaques, but infusion of a T-cell-depleting anti-CD8 β mAb in the

6 controller animals led to depletion of CD8⁺ T cells and the rapid reappearance of plasma viremia, suggesting that the prolonged suppression of viremia observed in these animals resulted from potent antiviral CD8⁺ T-cell immunity induced by the short course bnAb treatment amidst low levels of viral replication (53).




Figure 2-5 Establishment of controller status in bnAb-treated animals inoculated with SHIV_{AD8-EO} by the intrarectal route (a-f). Plasma viral loads (black) and CD4⁺ T-cell levels (red) are shown in 3 controller (a-c) and 3 noncontroller (d-f) macaques. From (53), Figure 2.




Figure 2-6 Establishment of controller status in bnAb-treated animals inoculated with SHIV_{AD8-EO} by the IV route (a-g). Plasma viral loads (black) and CD4⁺ T-cell levels (red) are shown in 3 controller (a-c) and 4 noncontroller (d-g) macaques. From (53), Figure 3.

3BNC117 and 10-1074 in combination, and 3BNC117 alone, have also been associated with virologic control in humans. Thirteen individuals with chronic HIV who received 2 or 4 3BNC117 infusions, immediately prior to analytical treatment interruption and then separated by 3 or 2 weeks, respectively, had a mean time to virologic rebound of 6.7 weeks (range of 5-9 weeks) after 2 infusions of the antibody or 9.9 weeks (range of 3-19 weeks) after 4 infusions of the antibody. Eleven chronically infected individuals who received 3BNC117 and 10-1074 immediately prior to and 3 and 6 weeks after analytical treatment interruption had a median time to virologic rebound of 21 weeks (range 5 to > 30 weeks) (54, 55) (see [Figure 2-7](#)).




Figure 2-7 Kaplan–Meier plots summarizing time to viral rebound for the participants with HIV-1 RNA < 20 copies per mL 2 weeks before and at the start of ATI (n = 11, left), for the participants sensitive to both antibodies (n = 9, center), and for the participants that showed pre-existing resistance to 1 of the antibodies (n = 2, right). The y axis shows the percentage of participants that maintain viral suppression. The x axis shows the time in weeks after start of ATI. Participants receiving the combination of 3BNC117 and 10-1074 are indicated by the blue line. Dotted red lines indicate a cohort of individuals receiving 3BNC117 alone during ATI (54) (n = 13) and dotted black lines indicate a cohort of participants who underwent ATI without intervention (8) (n = 52). From Mendoza et al (55), Figure 1c.

2.3.3 bnAbs modulate the autologous immune response to acute HIV infection

The mechanisms underlying bnAb-mediated virologic control are unknown, but there are clues in experiments that highlight bnAb-associated enhancement of autologous cellular and humoral responses. Adoptive transfer experiments have demonstrated that 3BNC117 can accelerate clearance of HIV-infected cells (see [Figure 2-8](#)) (56).




Figure 2-8 Percentage of Gag+ cells among CD3+CD8- cells in mice treated with 3BNC117 (600 mcg) or mice treated with isotype control 5 hours after HIV-1_{YU2}-infected cell transfer. From (56) Figure 2A.

Barouch et al demonstrated increased CD8⁺ T cell-mediated virus inhibition and decreased CD8⁺ and CD4⁺ T cell exhaustion in the presence of PGT121 compared to control as well as increased titers of autologous neutralizing antibody in macaques receiving PGT121, 3BNC117, and b12 (see [Figure 2-9](#) and [Figure 2-10](#)) (57).




Figure 2-9 SHIV_{SF162P3} and SHIV_{SF162P4} serum neutralizing antibody titers (ID₅₀) in the macaques described in [Figure 2-10](#) after administration of the triple PGT121, 3BNC117, and b12 monoclonal antibody cocktail. From (57) Extended Data Figure 2.




Figure 2-10 Therapeutic efficacy of the triple PGT121, 3BNC117, and b12 monoclonal antibody cocktail. LEFT: Log inhibition of viral replication in CD8⁺ T-lymphocyte-dependent virus suppression assays after monoclonal antibody infusion. One animal had no recoverable virus at week 2. CENTER and RIGHT: PD-1Ki671 expression on Gag-specific CD8⁺ (center) and CD4⁺ (right) T lymphocytes after monoclonal antibody infusion. From (57) Figure 1f, 1g, and 1h.

Enhanced humoral responses—specifically, elevated levels of HIV Env-specific antibodies—were also observed in macaques by Bolton et al (52). In this study, antibody titers in animals that received either VRC01 or VRC07-523LS + PGT121 with ART exceeded titers achieved by animals receiving ART alone (see [Figure 2-11](#)).




Figure 2-11 Geometric mean endpoint titer values for plasma reactivity to HIV-1 SF162 gp120 (anti-HIV Env) determined by enzyme-linked immunosorbent assay (ELISA) for each group (n = 6) at the indicated days postinfection. Error bars represent the 95% confidence intervals; the dotted lines indicate the limit of detection for the assay. From (52) Figure 3B.

Haigwood et al (58) demonstrated accelerated development of autologous neutralizing antibodies in macaques. Uninfected macaques were assigned to no immunization (n = 4) or passive immunization, 1 and 14 days after challenge, with either normal immunoglobulin G (IgG) (n = 6) or specific intravenous immunoglobulin (SIVIG) (n = 6), which is a polyclonal IgG obtained from a healthy SIV-infected nonprogressor animal with high-titer neutralizing activity. Three of the 6 animals that received SIVIG showed accelerated development of neutralizing antibodies, in contrast to the animals that received normal IgG or no IgG (see Figure 2-12). These SIVIG-associated antibodies contributed to rapid control of post-acute phase viremia, resulting in delayed disease onset.




Figure 2-12 Scatter plot of Env-specific binding and SIVsmE660 neutralizing antibodies (nAbs) at weeks 2 and 12 in macaques that seroconverted. Note that some symbols overlap. Mean ratios are noted for each group at weeks 2 and 12, with 95% confidence intervals. From (58), Figure 5.

Schoofs et al (59) also observed significantly increased autologous neutralizing antibody breadth and potency in chronically infected viremic individuals who received 3BNC117 compared to controls (see [Figure 2-13](#)).




Figure 2-13 Heterologous antibody responses. (A) The difference in overall area under the curve (AUC) (mean AUC change) per individual in TZM.bl assays against 13 heterologous viruses for d0 versus week 24 IgG obtained from 36 untreated viremic controls (mean sampling interval 26.8 weeks), 15 viremic individuals infused with 3BNC117 (mean sampling interval 24.1 weeks), and 12 ART-treated individuals receiving 3BNC117 infusion (mean sampling interval 24.0 weeks). (B) The aggregated differences in AUC between d0 and week 24 IgG assayed by TZM.bl for all viruses and all individuals. Each dot represents a single AUC difference for a single virus from one individual displayed in (A). Colored bars represent the mean of all AUCs. (C) 3BNC117 antibody levels (ELISA, white) and TZM.bl neutralization titer against tier 2 strain Q769.d22 (green) in patient 2A3. (D) Mean AUCs of IgGs of all individuals at d0 (gray) and week 24 (color of respective group) for each HIV-1 pseudovirus tested. Changes in neutralization of viremic control individuals without 3BNC117 infusion are shown in yellow (left). Change in neutralization of 3BNC117-treated

individuals are shown in dark blue (off ART, middle) and light blue (on ART, right). Red stars indicate significant p-values after Bonferroni correction (threshold $p < 0.0038$). From (59), Figure 2.

Furthermore, bnAbs appear to render otherwise-ineffective autologous antibodies more effective. In a hu-mice model in animals lacking autologous neutralizing antibodies, 10-1074 administered in combination with an autologous tier-1 neutralizing antibody (either 10-188 or 1-79) was associated with significantly greater reduction in viremia than 10-1074 or the autologous bnAbs alone. Additional experiments (data not shown) suggested that bnAbs directed viral evolution to variants more sensitive to autologous tier-1 nAbs, while the autologous nAbs also appeared to limit viral evolution away from the bnAb (see Figure 2-14) (60).




Figure 2-14 Changes of log₁₀ viral load (mean and standard error of the mean [SEM]) of mice treated with 10-1074 (gray), 10-1074 + 10-188 (blue), and 10-1074 + 1-79 (green). Significant differences between treatment groups were determined by using repeated measures analysis of variance (ANOVA) with a Bonferroni pos-hoc test. *, $p < 0.05$; **, $p < 0.001$. From (60) Figure 7D.

The concurrent enhancement of both cellular and humoral responses, and the associated evidence of virologic control, calls to mind the immunologic responses to tumor antigens in individuals receiving cancer immunotherapy, and implicates immune complexes formed by bnAbs bound to circulating virus and infected cells (61, 62). Immune complexes act as potent immunogens, as evidenced by work in the HIV vaccine field. A trial of an immune complex vaccine reported that anti-CD4bs complexes with gp120 led to conformational changes in gp120 that stabilized and exposed epitopes in the V3 loop, leading to enhanced antigenicity and immunogenicity compared to gp120 alone (63, 64). This recapitulates the difference in HIV acquisition without bnAbs present (ie, comparable to gp120 alone) compared to acquisition in the presence of bnAbs bound to the gp120, forming immune complexes that stimulate a more effective autologous immune response.

Immune complexes with infected cells and with cell-free virions undergo FcγR-mediated uptake by antigen presenting cells (eg, dendritic cells) and efficient antigen processing via proteasomal and endosomal degradation, that exposes particularly antigenic epitopes for MHCII presentation, stimulating enhanced CD4⁺ T-cell responses and directing the immunologic response down a Th1 pathway, with enhanced CD4⁺ co-stimulation, a more efficient cytokine milieu

for antiviral control, and enhanced CD8⁺ T-cell responses. The consequent development of pools of precursor T cells, in particular, sets the stage for durable virologic control, as these cells proliferate in the presence of antigen later, producing large amounts of effector T cells that then clear HIV-infected cells and virions (65-67). In fact, this pathway has been demonstrated in the murine FrCasE retrovirus model that has many similarities to HIV infection. T_{reg} suppression was also observed in this model after therapeutic bnAb administration and subsequent immune complex-mediated induction of an antiviral host response that preserved immune function and stymied disease development, demonstrating yet another potential contribution to the bnAb-optimized early antiviral responses that may ultimately yield long-term virologic control (68, 69).

Rendering this enhancement of the innate component of the immune response even more effective is bnAbs' binding and neutralization of virus—a direct “debulking” of viral particles and of infected cells, thus limiting the number of virions capable of direct infection and replication and serving to limit viral factors that impair dendritic cell activation and maturation. This “viral debulking” function of bnAbs is analogous to surgical debulking (ie, cytoreductive surgery) performed for some tumors, which is associated with a survival advantage that is attributed in part to improved autologous immune function, with lower levels of tumor factors that impair the host immune response to cancer.

In summary, by forming immune complexes, modulating and enhancing autologous immune responses, dampening HIV replication, and exerting a sieve effect at acquisition and pressure in early infection, even bnAb concentrations that are insufficient to prevent acquisition might nevertheless work in concert with the autologous innate and adaptive immune response to facilitate a more effective response to early infection and to help set the stage for later durable virologic control (70-72).

2.3.4 bnAbs may limit establishment and decrease the size of the viral reservoir

In addition to, and in part likely due to, their enhancement of innate and adaptive immune responses, bnAbs appear to be capable of limiting the establishment and decreasing the size of the viral reservoir (45, 73). An in-vitro study of cells obtained from chronically infected aviremic humans demonstrated bnAb binding of cell-free virions produced from the latent reservoir, suppression of entry of those virions to activated uninfected CD4⁺ T cells, and suppression of replication-competent HIV in autologous CD4⁺ T cells (74). In a humanized mouse model, administration of bnAbs 4 days after infection interfered with the establishment of a reservoir through Fc-mediated mechanisms and, when administered with latency-reversal agents, decreased the size of an established viral reservoir (40). Administration of bnAb with ART to reduce viral load, then of a single bnAb alone after ART termination, was followed by functional virologic control, demonstrating potential synergy between bnAbs and ART for durable viral suppression (75). In macaques, a reduction in cell-associated viral DNA and RNA

in PBMCs and tissues was also observed after administration of single and combination bnAb immunotherapy (57, 76).

The mechanisms of these impacts on the viral reservoir are likely myriad and include direct neutralization of virions from reactivated cells and bnAb binding and Fc-mediated recruitment of cytotoxic immune cells (eg, natural killer [NK] cells) that kill reactivated cells, which may be significantly enhanced by immune complex formation as described above (40, 45, 67). BnAbs accomplish this work of limiting reservoir seeding, and later reducing reservoir size, not only in blood and interstitial fluid but also in some of the principal sites of pre-ART virus production and persistence, lymph nodes and gut-associated lymphoid tissue, in which antiretrovirals have less penetrance (77-79). In one study, a significant reduction of cell-associated SHIV DNA was observed in these tissues after administration of a single bnAb to infected macaques (57) (see [Figure 2-15](#)).




Figure 2-15 Therapeutic efficacy of the single monoclonal antibodies PGT121 and 3BNC117. a–c, Plasma viral RNA (log copies/mL) in rhesus monkeys chronically infected with SHIV-SF162P3 after a single infusion (arrows) of PGT121 (a), 3BNC117 (b), or the control monoclonal antibody DEN3 (c). d–f, Proviral DNA (copies/10⁶ cells) in PBMCs (d), lymph nodes (e) and gastrointestinal mucosa (f) 14 days after the monoclonal antibody infusion in the animals that received PGT121 or DEN3. Red bars indicate mean values. Assays for one of the DEN3-treated controls failed. g, h, PD-1⁺ Ki67⁺ expression on Gag-specific CD8⁺ (g) and CD4⁺ (h) T lymphocytes in the monkeys that received PGT121. From (57), Figure 3.

2.4 Antiretroviral analytical treatment interruption (ATI)

More than 150 studies with ATIs have been conducted since the introduction of combination ART over 20 years ago. Nearly 100 of those were conducted in the first decade of the 21st century, largely seeking to mitigate the risk of ART toxicity, multiresistant virus, and treatment failures (80-85). Among these was the

Strategies for Management of Antiretroviral Therapy (SMART) trial, which evaluated standard continuous ART (n = 2,720) vs episodic, CD4+ T cell count-guided ART (n = 2,752). In 2006, the SMART team reported a hazard ratio (HR) of 2.6 (95% CI 1.9-3.7, p < 0.001) for the combined primary study endpoint of opportunistic disease or death from any cause when comparing episodic ART with standard continuous ART. Of note, the trial enrolled participants with CD4+ counts of > 350 and with any ART history, did not exclude participants with chronic hepatitis C virus (HCV) or hepatitis B virus (HBV) infection, monitored participants every 2 months in the first year and then every 4 months thereafter, and re-initiated ART in the episodic arm only when CD4+ T-cell counts dipped below 250 cells/mL. There were no virologic criteria for re-initiating ART, and the median duration of the first ATI in the episodic ART group was 16.8 months (81).

A subset analysis of the SMART data revealed that all serious adverse events (SAEs) in the trial occurred when CD4+ T cell counts were below 300 cells/mL. This subset analysis was restricted to 16 weeks of treatment interruption, participants with CD4+ T-cell counts > 400 cells/mL, ART re-initiation at CD4+ T-cell count of 350, and participants with no evidence of hepatitis B, hepatitis C, or diabetes at study entry. In participants meeting these criteria, no deaths and no renal or hepatic events were observed; 2 cardiac events were observed, one in each study group; and there was no difference in AIDS-related events (0.005% in the continuous ART group and 0.002% in the episodic treatment group) (86). These findings of no mortality or treatment-interruption-associated morbidity in this SMART subset are consistent with observations reported for several concurrent and subsequent treatment interruptions (all smaller than SMART; approximately half were randomized clinical trials like SMART) that similarly limited eligibility for and duration of treatment interruption, re-initiated ART at higher CD4+ cell counts, and monitored participants more frequently (84, 87-92).

Treatment interruption designs post-SMART, including the design for HVTN 805/HPTN 093, reflect lessons learned from SMART to mitigate the risks of treatment interruptions. Recent consensus guidelines on the design and conduct of analytical treatment interruptions further codify these risk mitigation measures and the current concept is aligned with these guidelines (see [Figure 2-16](#), (2, 3))

Panel: Key recommendations

Inclusion criteria

- Stable CD4 counts ≥ 500 cells per μL *
- HIV RNA undetectable on stable ART†
- Otherwise healthy individuals without major comorbidities

Key exclusion criteria

- Active or chronic hepatitis B virus infection, with detectable hepatitis B surface antigen, hepatitis B virus DNA, or both
- Active hepatitis C virus infection, with detectable virus RNA
- Active Mycobacterium tuberculosis infection‡
- History of systemic cancers, such as Kaposi's sarcoma and lymphoma, or other virus-associated malignancies§
- History of HIV-associated dementia or progressive multifocal leukoencephalopathy
- Resistance to two or more classes of antiretroviral drugs¶
- History of cardiovascular event or at high risk of an event (eg, atherosclerotic cardiovascular disease score $> 15\%$)
- History of AIDS-defining illness according to Centers for Disease Control and Prevention criteria
- History of CD4 nadir < 200 cells per μL during chronic stages of infection
- Women who are pregnant or breastfeeding
- Advanced non-alcoholic fatty liver and advanced nonalcoholic steatohepatitis, if evidence for substantial fibrosis (fibrosis score $\geq F2$) or evidence of cirrhosis
- HIV-related kidney disease or moderate-to-severe decrease in estimated glomerular filtration rate (< 45 – 60 mL/min/1.73 m^2)
- Children younger than 2 years of age when the ATI is planned

Monitoring

- HIV RNA monitoring weekly for 12 weeks, then every other week
- CD4 count monitoring every two weeks
- Monitoring of clinical symptoms, in particular in people who started ART during the hyperacute HIV phase
- Monitoring of participants' psychosocial experiences

ART restart criteria

- If requested by the participant or their HIV health-care provider
- If participant becomes pregnant
- If ART is deemed medically necessary for non-HIV related causes
- Symptomatic HIV disease||
- Confirmed absolute CD4 value < 350 cells per μL or CD4% $< 15\%$ **
- HIV RNA ≥ 1000 copies per mL for 4 weeks††
- Absolute HIV RNA $> 100\,000$ copies per mL††

Reducing risk of HIV transmission to sexual partners

- Offer pre-exposure prophylaxis and HIV testing referral information that trial participants can provide to their sexual partners

Additional or more stringent criteria might be required based on known toxicities of the study drug(s) or expected risks of the study intervention(s). Inclusion and exclusion criteria, monitoring, and antiretroviral therapy (ART) restart criteria might differ in children depending on age. ART=antiretroviral therapy. *Baseline CD4 counts of ≥ 350 cells per μL might be considered. †Based on FDA-approved HIV RNA quantification assay. ‡Latent tuberculosis infection discussed in the text. §Other malignancies discussed in the text. ¶Defined as single key mutations or an accumulation of minor mutations that result in resistance to entire respective drug classes. ||Symptoms include, but are not limited to, unintentional weight loss (> 5 – 10% of the pre-ATI bodyweight), otherwise unexplained persistent fever ($> 100.4^{\circ}\text{F}/38^{\circ}\text{C}$), persistent night sweats, persistent diarrhoea, oral candidiasis and generalised lymphadenopathy. **Largely dependent on the CD4 entry criteria; a sufficiently large delta between the entry value versus CD4 measurement for ART resumption should be ensured. ††12–16 weeks of uncontrolled viraemia, with HIV RNA of more than $100\,000$ copies per mL; it might be acceptable in studies in which a stable viral set point is a primary endpoint.

Figure 2-16 Consensus recommendations of the July 9, 2018 forum held at the Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, USA. From (3).

Specifically, the eligibility criteria recommended in Julg et al (3) and proposed by the HVTN 805/HPTN 093 team help ensure that the risks of hepatic, renal, and

cardiac morbidity and mortality observed in SMART are substantially reduced, as individuals are only eligible for treatment interruption if they have no active or chronic hepatitis B, no active hepatitis C, no evidence of other significant liver disease, no significant or unstable cardiac or cerebrovascular disease, and no evidence of HIV-related renal disease or moderate-to-severe reduction in estimated glomerular filtration rate (eGFR).

Eligibility and ART re-initiation criteria and the frequency and type of monitoring also reduce the risks of HIV-related morbidity and mortality observed in the SMART trial. These safety features include eligibility only after an extended period of virologic suppression on ART; requirement that volunteers on NNRTI-based ART regimens switch to short-acting antiretrovirals (Protease Inhibitor [PI]- or Integrase Strand Transfer Inhibitor [INSTI]-based ART regimens) in order to prevent emergence of ART-resistant HIV; requirement that volunteers have neither a history of AIDS-defining illness nor of specified HIV-associated conditions; eligibility only at higher baseline CD4⁺ T-cell counts; ART re-initiation for any symptomatic HIV disease, CD4⁺ T-cell counts < 350 cells/mL, or HIV RNA $\geq$ 1,000 copies/mL for 4 weeks; and initial monitoring weekly and then biweekly, including CD4⁺ T-cell counts, viral load, and psychosocial experiences.

The evidence that post-SMART changes in ATI trial design have, in fact, limited the risks of treatment interruption, is growing. For example, Huiting et al (93) demonstrated that among 22 participants who initiated ART in early HIV infection and then underwent a 16-week treatment interruption followed by re-initiation of ART, there were no significant differences between pre- and post-ATI CD4⁺ or CD8⁺ T-cell counts or percentage and no significant differences in reservoir markers, including HIV DNA, cell-associated RNA, and replication-competent HIV (see [Figure 2-17](#)).




Figure 2-17 Kinetics of HIV reservoirs and immunologic parameters prior to and following ATI and re-initiation of ART. The frequency of CD4+ T cells carrying HIV DNA and cell-associated viral RNA was measured prior to (pre-ATI) and following ATI (ATI) and upon re-initiation of ART (post-ATI). The frequency and cell count of CD4+ and CD8+ T cells are shown at the pre-ATI, ATI, and post-ATI timepoints. Statistical significance was tested with Wilcoxon signed rank test. *p < 0.05, **p < 0.01, ****p < 0.0001, NS, not significant. Adapted from (93) Figure 1.

Bar et al (48) reported on 24 chronically infected participants in 2 cohorts; in one cohort (A5340), participants had been taking ART for 3.8-6.0 years (median 4.7) and in the other (15-I-0140) participants had been taking ART for 7.7-13.3 years (median 10.0). All participants received VRC01, followed by an ATI, which ended upon confirmation of viral rebound. Median time to viral rebound was 4 weeks in the A5340 trial and 5.6 weeks in the 15-I-0140 trial. VRC01 slightly delayed plasma viral rebound compared with historical controls but did not sustain viral suppression. No safety issues were observed in either cohort.

Clarridge et al (94) reported on the 10 chronically infected participants in the 15-I-0140 cohort who received VRC01 immediately before and during ATI (median of 8 weeks, range 3-17 weeks). The length of the treatment interruption was governed by the time to meeting ART re-initiation criteria, which included a decrease of > 30% from the baseline CD4+ T-cell count or an absolute CD4+ T-cell count < 350 cells/mL, sustained (≥ 4 weeks) plasma viremia > 1,000 copies/mL, any HIV-related symptoms, or pregnancy (48, 94). During treatment interruption, total HIV DNA in CD4+ T cells increased significantly from pre-interruption levels and all participants rebounded, with median peak viremia of 30,950 copies/mL observed (range 340 - 273,221). By 6-12 months post-ATI, reservoir size, including the proportion of near full-length, genome-intact proviral sequences and structurally defective proviral sequences, and immune markers of exhaustion and activation returned to pre-ATI levels; CD4+, CD8+ (including activated CD8+), B, and NK cell levels also returned to pre-ATI levels; and there

was no evidence of emergence of antiretroviral drug resistance mutations in intact proviral DNA sequences.

Similar findings were reported by Salantes et al (95) among 14 chronically infected individuals who received VRC01 immediately prior to and during treatment interruption (median of 6 weeks, range 3-14 weeks) in A5340. The length of the treatment interruption was governed by the time to meeting ART re-initiation criteria, which included CD4⁺ T-cell count < 350 cells/mL or a return of HIV viremia defined as HIV RNA $\geq$ 200 copies/mL followed by HIV RNA $\geq$ 1,000 copies/mL or 3 consecutive measurements of $\geq$ 200 copies/mL. No significant changes in pre or postinterruption reservoir size or composition were observed by quantitative polymerase chain reaction (PCR), cell-associated RNA or quantitative viral outgrowth assay (QVOA). Perhaps most relevant to HVTN 805/HPTN 093, these findings remained true for participants with very small and relatively less heterologous reservoirs pre-ATI (eg, participant 7 in Figure 2-18).




Figure 2-18 Quantitative measures of reservoir change. Pre-ATI and post-ATI values (obtained at study entry and more than 6 months after viral suppression following ART re-initiation, respectively) of total HIV-1 DNA in CD4⁺ T cells (A), caRNA in CD4⁺ T cells (B), and the frequency of resting CD4⁺ T cells bearing replication-competent virus (infectious units per million [IUPM]) (C) are shown for each A5340 participant. Total DNA is not shown for participant A06, because the values were below detectable levels at both timepoints. P-values shown in A–C indicate the significance of the within-participant changes and were determined by Wilcoxon signed-rank test. From (95), Figure 2.

Strongin et al (96) assessed reservoir measures in 12 early- and chronic-treated participants across 4 ACTG studies that included ATI. Participants had at least 1 year of suppressive ART prior to treatment interruption that lasted a median of 12 weeks. No change in integrated HIV DNA was observed between baseline, pre-ATI samples and samples 6 months after ART re-initiation. In a shorter-term ATI, with a median duration of approximately 4 weeks, Papasavvas et al (97) demonstrated similar findings among 23 chronically-treated, ART-suppressed individuals. As expected, treatment interruption was accompanied by decreases in CD4⁺ T cells and increases in HIV viral load, markers of T-cell activation (eg, HLA-DR and CD38 co-expression), and levels of cellular HIV DNA (total DNA and 2-LTR [long terminal repeat] circles in PBMCs) and cellular RNA (by PCR). However, upon re-initiation of ART, viral re-suppression was achieved at a

median of 13 weeks and levels of CD4+ T cells, T-cell activation markers, and cellular HIV DNA and RNA returned to baseline, pre-ATI levels.

A recent meta-analysis offers further reassurance of the safety of ART treatment interruption with respect to rates of adverse events experienced by participants in trials with treatment interruptions. The meta-analysis included 7,104 participants across 22 studies with varying ATI designs, allowing for comparisons of outcomes across trials with different eligibility criteria, treatment interruption duration, follow-up intervals and ART re-initiation criteria. The overall rate of adverse experiences (AEs) during ATIs longer than 4 weeks' duration was 3% (95% CI 0%-7%) and was lower in studies with monitoring $\leq$ every 2 weeks (0%; 95% CI 0%-1%) than in studies with less frequent follow-up (6%; 95% CI 2%-13%; p-value for interaction = 0.01). Among the studies reporting new viral resistance mutations emerging during ATI, a 3% (95% CI 0%-8%) rate of such mutations was observed, with a higher rate in studies with less frequent monitoring (9%; 95% CI 3%-16%) compared to the rate observed in trials with more frequent ($\leq$ every 2 weeks) monitoring (0%; 95% CI 0%-1%; p = 0.03). Of note, the analysis was unable to discern which ART regimens preceded ATI in all of these trials, thus limiting their ability to account for trials with NNRTI switch periods to reduce the risk of the development of viral drug resistance mutations. Baseline CD4+ T-cell count was also associated with outcomes, with baseline CD4+ T cells > 500 cells/mcL associated with the fewest AEs (0%; 95% CI 0%-3%) (98, 99).

The authors conclude that treatment interruption can be conducted safely—with no lasting negative impact on clinical, immunologic, immune activation or virologic parameters—when appropriately designed with frequent monitoring and appropriate duration, eligibility and re-initiation criteria. The cited authors of the meta-analysis and assessments of ATI impacts described above specifically note that appropriate re-initiation criteria should allow for viral spikes (ie, without re-initiating ART prematurely) that may be necessary to elicit the ART-free, virologic suppression potential of immunomodulatory interventions, like bnAbs (18, 21, 93, 98-100).

The mitigation of risk in treatment interruptions relies on the optimal implementation of eligibility assessment, monitoring, timely ART re-initiation, and other risk mitigation practices. An assessment of site capacity to implement these aspects of the treatment interruption design—for example, capacity for weekly and biweekly clinic visits, including retention and adherence metrics thus far, and access to ART for timely re-initiation, including metrics of initial ART uptake—was completed prior to the determination of participating sites. Ongoing monitoring of the implementation of these risk mitigation measures will be in place throughout the trial's conduct. However, even with these measures, treatment interruptions do still entail risk and promise no anticipated benefit. These points must be made clear in an informed consent process for individuals contemplating any interruption in their antiretroviral therapy.

A pause in antiretroviral therapy—with careful eligibility criteria, frequent clinical and laboratory monitoring, and clear criteria for ART re-initiation—is a safe and acceptable strategy to evaluate the efficacy of VRC01 in this population of antiretroviral-treated HIV-infected adults. Current and prior studies of immune-based therapy, including a subgroup analysis of the SMART study, support the value, safety and acceptability of interruptions in antiretroviral therapy to assess virologic efficacy in this context (21, 48, 54, 86, 93, 101-105). In addition, while there remains uncertainty regarding how best to measure latent HIV reservoirs, several recent studies of participants undergoing similar short-term interruptions in antiretroviral therapy have noted no persistent changes in the size of the latent reservoir of infected CD4⁺ T cells following re-suppression by ART, based on the measures currently thought to reflect HIV reservoir size (21, 93-99). Specifically, ATI with intensive monitoring evaluating time to viral rebound has been advanced as a careful approach to provide insight regarding novel therapeutic approaches to control HIV off ART (79, 106, 107).

2.5 An AMP ATI

AMP participants who acquired HIV are unique: they were diagnosed with HIV soon after acquisition due to monthly HIV diagnostics on-study; they initiated ART soon after diagnosis due to AMP-facilitated linkage to ART; and those randomized to receive VRC01 had circulating bnAb at the time of acquisition due to the q8-week frequency of infusion and VRC01 PK. This sets the stage for formation of immune complexes of VRC01 bound to free virions or HIV-infected cells which, as detailed above, can modulate the immune response and, thus, the course of infection. This cohort offers a singular opportunity to assess the potential impact of very early immunotherapy on the immunologic and virologic trajectory of infection, a contribution that recent preclinical and clinical data suggests could meaningfully enhance our understanding of HIV pathogenesis and illuminate potential host defenses, and means of enhancing them, that previous HIV preventive, therapeutic, or curative interventions have heretofore not identified or successfully engaged.

A pause in antiretroviral treatment is the only way to determine if the presence of VRC01 at the time of, or shortly after, acquisition of HIV results in a blunted, delayed, or absent plasma viral rebound off ART, or subsequent control of rebound that may be observed. If such an impact on viral rebound is observed, ATI is also the best means of assessing mechanisms and predictive biomarkers of that impact through a detailed assessment of innate, cellular, and humoral markers along with characterization of the viral reservoir.

Recent research has suggested that viral rebound may be predicted by duration of infection, length of ART suppression, CD4⁺ T-cell nadir, total HIV DNA and cell-associated HIV RNA, among others; but samples and validation of these potential predictors are limited and there is no consensus biomarker predictive of durable viral suppression (8, 77, 108). Identification of reliably predictive

biomarkers could inform future therapeutic and cure trial design and guide allocation of limited resources, advancing promising therapies more efficiently, and limiting exposure of trial participants to ineffective interventions. The AMP trial has pre and postinfection serum and limited PBMC samples on each participant, including samples obtained ≤ 4 weeks from the time of acquisition and pre-ART. Thus, across currently available and proposed samples, ATI among AMP participants provides a unique opportunity to contribute to the exploration and validation of biomarkers, many observed in the preclinical studies described in Section 2.3.2 through 2.3.4 above, that may predict the virologic and immunologic course of infection among bnAb and placebo recipients.

2.6 Ethical considerations

In the last 2 decades of treatment interruption conduct, an ethical framework for the conduct of such studies has evolved. There is general agreement that the risks of treatment interruption must be minimized; the trial must provide valuable data to answer an important question; and there must be no other methods entailing less risk that could answer that question (109, 110).

While ethicists have noted that “in general, the risks of the ATI itself, when closely monitored, are considered to be low,” risk mitigation is an essential part of the design of any treatment interruption. Risk mitigation includes but is not limited to: adoption of appropriate eligibility criteria; efforts to enhance potential participant benefits; community engagement in the design and implementation of the trial; frequent participant monitoring, balanced with considerations of participant burden; and a sound informed consent process. The informed consent process should acknowledge the risks and the lack of anticipated individual benefits, the fact that treatment interruptions are not recommended in clinical care, and the fact that ART guidelines recommend lifelong treatment (109, 110).

Each of these recommendations has been adopted by the HVTN 805/HPTN 093 Protocol Team. The HVTN 805/HPTN 093 informed consent process and Sample Informed Consent Form (SICF) include all points noted above and include elements demonstrated to improve the quality of informed consent in not only HIV research but also other fields (see Section 12.1). Trial design, including eligibility and ART re-initiation criteria, ATI duration, and monitoring frequency, are modeled on the most recent consensus guidelines for treatment interruption design (see Figure 2-16 above) (2, 3). The HVTN 805/HPTN 093 Protocol Team will implement not only rigorous counseling for participants on reducing HIV transmission risk to their partners but also a program for PrEP provision to participants’ partners while the participant is enrolled; HIV genotyping (not standard of care in many AMP regions, unless first-line ART options have been exhausted); and accelerated, physician-led ART (re-initiation) access for participants in regions where ART stock-outs, public nurse-led care, and a window of several weeks between meeting ART re-initiation criteria and ART initiation could pose challenges for participants.

As all potential participants in HVTN 805/HPTN 093 would come from the parent HVTN 703/HPTN 081 study, early engagement of investigators and community representatives in several communities where HVTN 703/HPTN 081 was conducted has already been initiated and is ongoing. These consultations with community stakeholders and investigators include an overview of the history and current context of treatment interruption, including potential risks and region-specific considerations. They also provide an opportunity for dialogue about the immunologic hypotheses and design for HVTN 805/HPTN 093 and began laying a foundation of authentic stakeholder engagement in the earliest weeks of development of this study. See Section 2.7 below for further details.

2.7 Feasibility

The HVTN 805/HPTN 093 team has the significant advantage of data and experience from the AMP Study to inform an assessment of the feasibility of HVTN 805/HPTN 093, appreciating that while this concept is significantly different from the parent AMP Study, all participants for HVTN 805/HPTN 093 will be from the AMP cohort at sites and in communities now familiar to the AMP and HVTN 805/HPTN 093 teams. AMP accrual and retention rates, early termination rates, data quality and safety have been excellent by any standard, perhaps particularly so considering the demanding nature of the 2 AMP trials (HVTN 704/HPTN 085 and HVTN 703/HPTN 081) that, together, enrolled 4,625 participants for 2 years of monthly clinic visits and q8-weekly IV infusions (111, 112). In over 35,000 IV infusions administered to date, no procedural complications (eg, local IV site infection,) have been observed; 2/3 of participants receive VRC01 and very low rates of infusion reactions have been observed, all of which have been medically managed exceptionally well (111, 113).

While overall performance in AMP has been exceptional, the HVTN 805/HPTN 093 team also considered site-specific performance and other considerations in determining which sites will implement HVTN 805/HPTN 093. Considerations include: site-specific retention performance; timeliness and accuracy of safety data submission; access to potentially eligible participants; site capacity for frequent clinic visits and close clinical monitoring of participants; access to and capacity for close partnership with primary HIV care, including timely ART re-initiation; and site willingness to participate, including receptivity of site investigators, community representatives, and regulatory authorities.

Partnership with regional investigator, regulatory, and community representatives for the development of HVTN 805/HPTN 093 has already begun in earnest and is ongoing.

A symposium with sub-Saharan African investigators, regulatory and public health representatives, and community representatives, including representatives from local Community Advisory Boards (CABs), was held in August 2019 in Cape Town, South Africa, with co-sponsorship by the South African Medical

Research Council (SAMRC). Attendees were engaged and were supportive of contributing to the global HIV cure and remission research agenda, including with rigorously-designed ATIs like HVTN 805/HPTN 093, with some participants stating, “*We need this,*” “*This is for our communities,*” and “*This [an AMP ATI] is a first step.*” Understanding and engagement was also evident in attendee questions and dialogue throughout the consultation.

Early in development in early 2019, the HVTN 805/HPTN 093 team also solicited and received input and support from sub-Saharan African AMP investigators, policy makers, basic and clinical scientists, and from the AMP Community Working Group (AMP CWG), whose members include community educators, recruiters, and CAB members from each AMP site. Several forums throughout 2019 also provided opportunities for additional discussion and feedback regarding HVTN 805/HPTN 093 with AMP Protocol Team and site representatives (eg, investigators, clinic coordinators) and AMP CWG members.

Community engagement and site readiness, as described above, are necessary but not sufficient to support the conduct of HVTN 805/HPTN 093. AMP participant willingness to participate in the ATI is a crucial additional feasibility consideration. Again, interim data from AMP provides some insight here.

A mixed-methods AMP sub-study enrolled 300 participants—50 AMP participants from each of 6 sites based throughout the US. Upon exiting the AMP Study, each participant took part in a brief survey and a 60-minute interview regarding their demographics, understanding of bnAbs, and any enrollment or retention barriers and facilitators they experienced. In an interim analysis, about a quarter of participants expressed concerns about the AMP Study time commitment or the experimental nature of the VRC01 product and the associated uncertainty regarding its efficacy. All participants expressed a strong sense of appreciation for and positive identification with their clinical research site. For example, one participant noted: “*I’m going to volunteer for clinical research, especially if it’s research that’s going to affect the community that I’m part of*” and another enthused “*I’m such a big fan of [the CRS] that any research study that they pretty much put forward, I’m like interested in, or definitely highly support.*” Several participants cited the additional medical care they received at the clinic as a facilitator of ongoing study participation: “*[staff member name] took on this role of like a secondary primary care physician for me. In addition to the care I was getting at my doctor’s office, I also felt like I was getting another set of eyes on me. I just felt cared for.*” Another observed “*I liked how I got tested so often, and they were there to help me with any concerns I had, whether in regard to the infusion or to getting tested for other STDs or STIs.*”

While the scope of the sub-study is limited to the US, anecdotal evidence from sites around the world, coupled with the exceptional accrual and retention observed in the AMP studies to date (112), suggests that the experience participants have with their HVTN and HPTN AMP sites, globally, is a strongly positive motivator for continued engagement with their site. Furthermore, many

sites retain engagement with their participants through various means even post-AMP and have the ability, prespecified in site-specific AMP consent forms approved by their Institutional Review Boards/Ethics Committees (IRBs/ECs), to reach out to participants after they've exited the trial to inquire about their interest in possible future research.

Thus, HVTN 805/HPTN 093 as described herein is thought to be feasible thanks in large part to the selection of sites for AMP and their conduct of the study, including their relationships with site participants and their community and regulatory partners.

2.8 Conclusion

The AMP ATI is uniquely positioned to test for ART-free virologic control and to enhance understanding of the immunologic (ie, host) and virologic mechanisms and biomarkers of control among individuals who acquired HIV in the presence of circulating bnAb. Here, we have an opportunity to translate the intriguing preclinical observations in SHIV-infected primates (53) to humans. This trial will extend early clinical observations, explore potential mechanisms, validate biomarkers of control, and discern whether AMP participants who became HIV-infected while on trial may become “post-bnAb controllers,” capable of early ART- and/or bnAb-induced immunologic responses that change the course of their infection.

3 Study design

3.1 Hypotheses

3.1.1 Primary hypotheses

- Individuals who initiated ART early in HIV infection in HVTN 703/HPTN 081 will suppress plasma viremia and maintain CD4+ T-cell counts longer during ATI than historical cohorts (ie, SPARTAC African and non-African cohorts described in Gossez et al (1) comprising individuals who initiated ART early in HIV infection without other interventions).
- Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will suppress plasma viremia and maintain CD4+ T-cell counts longer during ATI than those who initiated ART early in HIV infection and received placebo within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081.
- ATI will be safe and well-tolerated in individuals who acquired HIV during their participation in HVTN 703/HPTN 081.

3.1.2 Secondary hypotheses

- Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will have enhanced cellular and humoral responses compared with those who initiated ART early in HIV infection and received placebo within 8 weeks of acquiring HIV in HVTN 703/HPTN 081.
- Individuals who initiated ART early in HIV infection and also received VRC01 within 8 weeks before or after acquiring HIV in HVTN 703/HPTN 081 will have more limited viral reservoirs, before and after ATI, than those who initiated ART early in HIV infection and received placebo within 8 weeks of acquiring HIV in HVTN 703/HPTN 081.

3.2 Objectives and endpoints

3.2.1 Primary objectives and endpoints

Primary objective 1

- To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early

infection on the time to meeting ART re-initiation criteria in participants undergoing ATI

Primary endpoint 1

- Time to meeting criteria for ART re-initiation
- Frequency of sustained post-treatment HIV control, defined as ≥ 24 weeks off ART without meeting ART re-initiation criteria

Primary objective 2

- To evaluate the safety of ATI among HVTN 805/HPTN 093 participants

Primary endpoint 2

- Laboratory measures of safety, adverse events (AEs), SAEs, and rates of discontinuation

3.2.2 Secondary objectives and endpoints

Secondary objective 1

- To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection on the development of anti-HIV immune responses and on the potential association of those immune responses with time to meeting criteria for ART re-initiation in participants undergoing ATI

Secondary endpoints 1

- Response rate, magnitude, and polyfunctionality of HIV-specific CD4+ and CD8+ T-cell responses as measured by flow cytometry
- Magnitude and breadth of neutralizing antibody (nAb) responses against autologous and heterologous HIV isolates, as measured by TZM-bl neutralization assay
- Non-neutralizing, FcγR-mediated antibody effector functions such as antibody-dependent cellular cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), and virion capture
- Frequency of dendritic cell activation and maturation markers, as measured by flow cytometry or other cell phenotyping assays
- Frequency of T- and B-cell activation and exhaustion markers, as measured by flow cytometry or other cell phenotyping assays

Secondary objective 2

- To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection on viral load in participants undergoing ATI

Secondary endpoint 2

- Cumulative incidence of participants with viral load ≥ 200 at weeks 8, 16, and 24

Secondary objective 3

- To evaluate the effect of early ART initiation, with or without having received VRC01 in the immediate pre-HIV acquisition period and/or during early infection on HIV reservoir size before and after ATI, and whether HIV reservoir measurements are associated with time to meeting criteria for ART re-initiation in participants undergoing ATI

Secondary endpoint 3

- Frequency of CD4+ T cells carrying intact and/or total pro-viral HIV DNA, replication competent virus, and/or cell-associated HIV RNA

3.2.3 Exploratory objectives

Exploratory objective 1

- To assess participant and potential participant motivations, perceptions and tolerance for ATIs, for prolonged viremia, and for trial requirements (eg, ART switches, barrier protection for all sexual activity)

Exploratory objective 2

- To assess social benefits (eg, altruism) and negative social impacts, including but not limited to anxiety regarding being off ART and becoming viremic again and fear of inadvertent HIV transmission to partners

Exploratory objective 3

- To assess participant and potential participant decision-making quality, including perception of being informed, clarity and support in decision making and satisfaction with their choice regarding participation

Exploratory objective 4

- To evaluate the effect of VRC01 received in the immediate pre-HIV acquisition period and/or during early infection on viral kinetics in participants undergoing ATI

Exploratory objective 5

- To determine whether characteristics of the transmitted/founder (T/F) viruses, rebound virus(es), and/or baseline viral reservoirs, including relative VRC01 sensitivity or resistance, are associated with virologic control or the development of anti-HIV immunity among individuals who acquired HIV during their participation in HVTN 703/HPTN 081

Exploratory objective 6

- To assess whether host genetics (eg, HLA type, CCR5 heterozygosity) and sex assigned at birth are associated with time to meeting ART re-initiation criteria among VRC01 or placebo recipients who acquired HIV during their participation in HVTN 703/HPTN 081

Exploratory objective 7

- To qualitatively characterize the immune responses and reservoir characteristics of controllers and non-controllers in this cohort and to compare these to immune responses and reservoir characteristics of historical cohorts (ie, SPARTAC African and non-African cohorts, described in Gossez et al (1), comprising individuals who initiated ART early in HIV infection without other interventions)

3.3 ATI

For all participants, ATI begins with cessation of ART in Schedule 1, *Monitoring ATI*. Participants on ATI (ie, Schedule 1 or 2) who do not meet criteria for moving to the next visit schedule (ie, viral rebound or re-initiating ART) should continue on their current ATI visit schedule, including extended visits (as appropriate) until such time as they move to the next visit schedule or are terminated from the study. Schedule 1 and Schedule 2 extended visits occur quarterly and are designated as Visits Type A and Type B, with procedures specified for each visit type, in the laboratory and clinical procedures tables (see [Appendix E](#), [Appendix F](#), [Appendix H](#), [Appendix I](#), and [Appendix K](#)). For participants who meet criteria for reinitiating ART but decline to do so, see Section [3.3.4](#).

3.3.1 Transition from Schedule 1, *Monitoring ATI* to Schedule 2, *ATI monitoring with viremia*

For participants on Schedule 1, *Monitoring ATI*, the following virologic criterion will trigger a transition to Schedule 2, *ATI monitoring with viremia*:

- Confirmed VL ≥ 200 copies/mL (ie, on 2 consecutive samples).

A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first VL result ≥ 200 copies/mL (see HVTN 805/HPTN 093 SSP for details).

3.3.2 Transition from Schedule 1, *Monitoring ATI* to Schedule 3, *Follow-up on ART*

For participants on Schedule 1, *Monitoring ATI*, any of the following non-virologic criteria will trigger ART re-initiation and transition to Schedule 3, *Follow-up on ART*:

- Confirmed CD4+ T-cell count < 350 cells/mm³ (ie, on 2 consecutive samples)
- Any HIV-related syndrome (eg, ARS, an opportunistic infection)
- Pregnancy or breastfeeding
- ART re-initiation requested by participant
- ART re-initiation requested by primary HIV care provider or clinical research site (CRS) Investigator of Record (IoR) (eg, if deemed medically necessary)

A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see HVTN 805/HPTN 093 SSP for details).

3.3.3 Transition from Schedule 2, *ATI monitoring with viremia*, to Schedule 3, *Follow-up on ART*

For participants on Schedule 2, *Monitoring ATI with viremia*, the following virologic criteria will trigger ART re-initiation and transition to Schedule 3, *Follow-up on ART*.

From the Week 0 visit to the Week 24 visit in Schedule 2:

- Viral load remains $\geq 1,000$ copies/mL for ≥ 4 consecutive weeks AND viral load has not dropped 0.5 log from the previous week

This ART re-initiation criterion applies irrespective of missed visits and must take account of VL results $\geq 1,000$ copies/mL experienced by a participant while on Schedule 1. Furthermore, assessing whether participants meet this criterion may require additional visits beyond those scheduled from Week 8 (Visit 48) to Week 24 (Visit 56) in Schedule 2. Ideally, these additional visits will be conducted weekly. These additional visits will be treated as interim visits (for more details, see the HVTN 805/HPTN 093 SSP).

After the Week 24 visit in Schedule 2, the ART re-initiation criterion is:

- Confirmed viral load ≥ 200 copies/mL (ie, on 2 consecutive samples).

A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first VL result ≥ 200 copies/mL (see HVTN 805/HPTN 093 SSP for details).

The above virologic ART re-initiation criteria are shown schematically in [Figure 3-1](#).



Figure 3-1 Virologic criteria for transition from Schedule 2 to Schedule 3 (ie, ART re-initiation)

In addition, at any time during Schedule 2, any of the following non-virologic criteria will trigger prompt ART re-initiation and transition to Schedule 3, *Follow-up on ART*:

- Confirmed CD4+ T-cell count < 350 cells/mm³ (ie, on 2 consecutive samples)
- Any HIV-related syndrome (eg, ARS, an opportunistic infection)
- Pregnancy or breastfeeding
- ART re-initiation requested by participant
- ART re-initiation requested by primary HIV care provider or CRS IoR (eg, if deemed medically necessary).

A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see HVTN 805/HPTN 093 SSP for details).

3.3.4 Participants who decline to re-initiate ART

When ART re-initiation criteria are met, CRS staff will strongly recommend that the participant restart ART. Participants who decline to do so should be urged to continue clinic visits per the schedule they are in at that time (eg, Schedule 1 or Schedule 2) until such time as the participant re-initiates ART, at which time they should transition to Schedule 3. Monitoring such participants may require additional visits. Ideally, these visits will be conducted weekly. These additional visits will be treated as interim visits. For details, see the HVTN 805/HPTN 093 SSP.

4 Statistical considerations

4.1 Estimated accrual

Recruitment will target participants who received VRC01 or placebo and became HIV-infected during participation in HVTN 703/HPTN 081.

We calculate the expected trial size under different scenarios for the percentage of enrolled participants out of the total number of endpoint infected cases from HVTN 703/HPTN 081; scenarios include 25%, 50%, and 75%. Given an assumption of a common prevention efficacy at each dose (PE10 and PE30), the expected trial size will depend on different levels of PE. Under the null hypothesis (PE10 = PE30 = 0%), the total expected number of endpoint infections among placebo recipients is 26 and VRC01 recipients is 54 (Table 4-3 in the parent protocol HVTN 703/HPTN 081); therefore, we assume a trial size of 7, 13, or 20 placebo recipients and 14, 27, or 41 VRC01 recipients depending on the enrollment ratio. Under the alternative hypothesis (PE10 = PE30 = 60%), the total expected number of endpoint infections among placebo recipients is 35 and VRC01 recipients is 28 (Table 4-3 in the parent protocol); therefore, under the different enrollment ratio scenarios, we assume a trial size of 9, 18, or 26 placebo recipients and 7, 14, or 21 VRC01 recipients. These 6 sample size scenarios represent estimated upper and lower bounds for sample size given the expected number of endpoint infections under the null and alternative hypotheses respectively, as well as various scenarios for the percentage of eligible participants who will ultimately enroll in HVTN 805/HPTN 093.

4.1.1 Power to evaluate time to ART re-initiation criteria

Using data from the placebo arm of a therapeutic vaccine study which enrolled early treated patients (21), we modeled the time T (in weeks) to meet re-initiation criteria as a Weibull distribution with survival function $P(T > t) = \exp[-(\lambda t)^p]$, where $\lambda = \exp(-2.932) = 0.053$ and $p = 2.014$. From these parameters, we estimated the median time to meet re-initiation criteria as 15.6 weeks in the AMP placebo recipients. We can express the treatment effect in terms of either the HR θ or the factor β by which treatment increases the median time to meet re-initiation criteria; β and θ are related by $\beta = \theta^{1/p}$.

We model censoring of time to restart criteria using an exponential model and a maximum follow-up time of 72 weeks, where follow-up is censored at a rate of 20% per year due to either study termination or re-initiation of ART before reaching the restart criteria. Power for a 40%, 50%, 60%, and 70% reduction in hazard under different prevention efficacy scenarios are shown in [Table 4-1](#). These hazard reductions correspond to 29%, 41%, 58%, and 82% increases in the median time to meet restart criteria. Therefore, if VRC01 increases the median time to restart criteria 58% and the enrollment rate is 50%, then under the parent

study null hypothesis power to detect a difference in time to restart criteria is 72% while under the alternative hypothesis power is 62%.

Table 4-1 Power under different prevention efficacy results, different enrollment rates and different treatment effects, expressed as either reduction in hazard or percent increase in median time to meet re-initiation criteria

Enrollment Rate		Null hypothesis			Alternative hypothesis		
		25%	50%	75%	25%	50%	75%
Expected number of placebos		7	13	20	9	18	26
Expected number of VRC01		14	27	41	7	14	21
Reduction in hazard	% Increase in median time to meet re-initiation criteria						
40%	29%	20%	31%	44%	16%	27%	36%
50%	41%	31%	49%	67%	23%	42%	57%
60%	58%	46%	72%	89%	35%	62%	81%
70%	82%	65%	90%	98%	52%	85%	95%

4.1.2 Sample size considerations for safety

The goal of the safety evaluation for this study is to identify safety concerns associated with ATI. The ability of the study to detect SAEs (see Section 4.2.3.1) can be expressed by the true event rate above which at least 1 SAE would likely be observed and the true event rate below which no events would likely be observed. Under the enrollment rate and PE scenarios described in Section 4.1, Table 4-2 shows the true event rate such that there is a 90% chance of observing at least 1 event (or no events) among three groups; 1) the placebo recipients only, 2) the VRC01 recipients only, or 3) all enrolled participants.

Table 4-2 True event rates above which at least 1 SAE (or below which no events) would likely be observed in a group of size n participants. Event rates are computed separately for various values of n given by the range and midpoint for the estimated number of participants among 3 groups; 1) the placebo recipients only, 2) the VRC01 recipients only, or 3) all enrolled participant based on the enrollment scenarios described in Section 4.1.

n	Placebo recipients			VRC01 recipients			All enrolled		
	7	17	26	7	24	41	16	39	61
At least 1 event	28.0%	12.7%	8.5%	28.0%	9.1%	5.5%	13.4%	5.7%	3.7%
No events	1.5%	0.6%	0.4%	1.5%	0.4%	0.3%	0.7%	0.3%	0.2%

4.2 Statistical analyses

Analyses for primary endpoints will be performed using SAS and R. All other descriptive and inferential statistical analyses will be performed using SAS, StatXact, or R statistical software.

No formal multiple comparison adjustments will be employed for multiple safety endpoints or secondary endpoints. However, multiplicity adjustments will be made for certain laboratory assays, as discussed below, when the assay endpoint is viewed as a collection of hypotheses (eg, testing multiple peptide pools to determine a positive response).

4.2.1 Analysis variables

The analysis variables consist of baseline participant characteristics, safety, and laboratory outcomes for primary- and secondary-objective analyses.

4.2.2 Baseline comparability

Groups defined by the parent protocol treatment arms will be compared for baseline participant characteristics using descriptive statistics.

4.2.3 Primary virologic analysis

In this study we either compare placebo recipients from HVTN 703/HPTN 081 with historical controls (ie, SPARTAC cohorts described in Gossez et al (1)) or we condition on HIV-infection with the intention of comparing virologic outcomes by randomized treatment assignment from HVTN 703/HPTN 081.

When comparing outcomes by treatment assignment among infected AMP participants, we are conditioning on a postrandomization event, HIV-infection. Consequently, a two-sample test for a difference in outcome is subject to postrandomization selection bias. A direct comparison of virologic outcomes between treatment groups, which measures the “net treatment effect,” does not have a causal interpretation (114). As an example of the type of bias that can occur, suppose VRC01 protects against mild viruses but is not effective against virulent viruses. If this were true, we might see longer time to re-initiation criteria in VRC01 recipients than placebo recipients whereas, if we restricted only to those participants infected with a virulent HIV strain, we might see shorter time to re-initiation criteria in the VRC01 group. Therefore, a two-sample test comparing time to re-initiation criteria between treatment groups could give a misleading impression that VRC01 shortens viral suppression time during ATI while the causal interpretation of this hypothetical example would be that for virulent viruses, VRC01 increases viral suppression time during ATI compared to placebo.

Approaches will be taken to ensure robust and unbiased results. Implementation of targeted minimum loss-based estimation (TMLE) methods, used to address the objectives in the following sections, will be fully prespecified in the statistical analysis plan (SAP) to ensure objective and reproducible inference.

When comparing by treatment assignment among AMP cases, each analysis will be done pooling the VRC01 recipients versus placebo recipients with

supplemental analyses done separately for each dose group versus the placebo recipients.

4.2.3.1 Time to ART re-initiation criteria

The primary outcome is the time from the start of treatment interruption until the participant meets criteria to re-initiate ART.

For a comparison of the placebo group to historical controls a Cox proportional hazards model with the indicator of AMP placebo group versus historical control group will be used to estimate the cumulative incidence in each group, as well as the HR, all with 95% confidence intervals and associated 2-sided p-values, where the model controls for baseline covariates thought to potentially predict both HIV-1 infection and the instantaneous hazard of meeting the ART re-initiation criteria. This is needed for controlling for potential selection bias given that analyzed treatment groups are not randomized. Plots of the estimated cumulative incidence will be shown by treatment group.

A Cox proportional hazards model with the indicator of assignment to a mAb group versus the control group will be used to estimate the cumulative incidence in each of the infected VRC01 and infected placebo groups, as well as the HR, all with 95% confidence intervals and associated 2-sided p-values, where the model controls for baseline covariates thought to potentially predict both HIV-1 infection and the instantaneous hazard of meeting the ART re-initiation criteria. This is needed for controlling for potential postrandomization selection bias given that analyzed treatment groups are selected postrandomization. Plots of the estimated cumulative incidence will be shown by treatment group.

As a supportive analysis of this hypothesis, TMLE may be used to estimate cumulative incidences of the primary efficacy endpoint over time for the pooled mAb arm and the control arm. Iterative mean-based TMLE is used for this analysis as described by Benkeser et al (115). The Super Learner (116) is used to generate initial estimates of the conditional censoring distribution and the iterated conditional means. This analysis will use TMLE as implemented in the R package *survtmle* available on CRAN (117).

4.2.3.2 Frequency of sustained post-treatment HIV control

The same covariate adjusted model used to estimate cumulative incidence of meeting the re-initiation criteria for ART over time, described in the previous section, will be used to compare the rates of failure to maintain HIV control at 24 weeks between the placebo group and historical controls and between the 2 treatment groups.

4.2.4 Secondary analyses of immune responses, reservoir measurements and viral load

Analyses of immune responses and reservoir measurements will be descriptive using appropriate plotting techniques for describing measurements at a fixed time point (eg, boxplots and barplots) or longitudinally (eg, spaghetti plots) separately by treatment group. In addition, if enough samples are assayed we plan to do inferential and descriptive analyses as described in the following sections.

4.2.4.1 Evaluate the difference in anti-HIV immune responses or reservoir size between VRC01 and placebo recipients

TMLE will be used to estimate mean endpoints in each of the infected VRC01 and infected placebo groups, as well as the mean difference, all with 95% confidence intervals and associated 2-sided p-values, where the TMLE controls for all baseline covariates thought to potentially predict both HIV-1 infection and one of the secondary endpoints under study. This is needed for controlling for potential postrandomization selection bias given that analyzed treatment groups are selected postrandomization.

4.2.4.2 Assess whether anti-HIV immune responses or reservoir size are associated with time to meeting ART re-initiation criteria

Cox regression will be used to estimate cumulative incidence over time for groups defined by levels of immune response or reservoir size for each treatment group.

4.2.4.3 Time to viral load > 200

Secondary endpoint 2 is time until the participant has a viral load > 200. The same methodology used to assess time to ART re-initiation criteria will be used for this endpoint. In this analysis we will compare cumulative incidence at week 8 in each of the infected VRC01 and infected placebo groups supplemented by a sensitivity analysis based on cumulative incidence at weeks 8, 16, and 24.

4.2.5 Primary safety analysis

4.2.5.1 AEs and SAEs

AEs will be summarized using Medical Dictionary for Regulatory Activities (MedDRA) System Organ Class and preferred terms. Tables will show by treatment arm the number and percentage of participants experiencing an AE within a System Organ Class or within preferred term category by severity or by relationship to treatment interruption. For the calculations in these tables, a participant with multiple AEs within a category will be counted once under the maximum severity or the strongest recorded causal relationship to treatment interruption. Formal statistical testing comparing arms is not planned since interpretation of differences must rely heavily upon clinical judgment.

4.2.5.2 Local laboratory values

Box plots of local laboratory values will be generated for baseline values and for values measured during the course of the study by treatment arm and visit. Each box plot will show the first quartile, the median, and the third quartile. Outliers (values outside the box plot) will also be plotted. If appropriate, horizontal lines representing boundaries for abnormal values will be plotted.

For each local laboratory measure, summary statistics will be presented by treatment arm and timepoint, as well as changes from baseline for postenrollment values. In addition, the number (percentage) of participants with local laboratory values recorded as meeting Grade 2 AE criteria or above as specified in the Division of AIDS Table for Grading the Severity of Adult and Pediatric Adverse Events will be tabulated by treatment arm for each timepoint after initiation of treatment interruption. Reportable clinical laboratory abnormalities without an associated clinical diagnosis will also be included in the tabulation of AEs described above.

4.2.5.3 Reasons for discontinuation of ATI and early study termination

The number and percentage of participants who discontinue ATI and who terminate from the study early will be tabulated by reason and treatment arm.

5 Study population

Participants for this trial will be recruited from among former HVTN 703/HPTN 081 (NCT02568215) study participants who met criteria for transition to Schedule 2 or Schedule 3 in that study and who meet the following inclusion/exclusion criteria.

Investigators should always use good clinical judgment in considering a volunteer's overall fitness for trial participation. Some volunteers may not be appropriate for enrollment even if they meet all inclusion/exclusion criteria. Medical, psychiatric, occupational, or other conditions may make evaluation of safety difficult, or may make volunteers unable to comply with the requirements of participation. For these reasons, some volunteers may be poor candidates for retention.

5.1 Inclusion criteria

1. **Estimated date of HIV-1 acquisition** within 8 weeks of (ie, before or after) having received an HVTN 703/HPTN 081 infusion (see HVTN 805/HPTN 093 SSP)
2. **Initiated ART within 28 weeks** of HVTN 703/HPTN 081 date of HIV-1 diagnosis
3. **Receiving continuous ART for at least 1 year**
 - ART interruptions of up to 7 days in duration and ≥ 90 days prior to enrollment are acceptable.
 - Within- and between-class changes in ART within the previous year are acceptable.
4. If on an NNRTI, **willingness and ability to switch** to a PI- or INSTI-containing regimen for at least 4 weeks prior to ART interruption
5. **Willingness to interrupt ART** for up to 24 weeks or up to the time of meeting ART re-initiation criteria (Section [3.3](#))
6. **Willingness to re-initiate ART** upon meeting study ART re-initiation criteria
7. **Willingness to use barrier protection** (ie, male or female condoms) for all sexual activity during the ATI and until confirmation of viral suppression following ART re-initiation
8. **Willingness for CRS staff to contact primary HIV care provider** to exchange information regarding HVTN 805/HPTN 093 and participant medical history

9. Site investigator anticipates that a **fully active alternative ART regimen** could be constructed and would be available in the event of virologic failure on the participant's current ART regimen.
10. **Access to a participating CRS** and willingness to adhere to study visit schedule and to be followed for the planned duration of the study
11. Ability and willingness to provide **informed consent**
12. **Assessment of understanding:** volunteer demonstrates understanding of this study; completes a questionnaire prior to enrollment with verbal demonstration of understanding of all questionnaire items answered incorrectly.
13. **Agrees not to enroll in another study** of an investigational research agent for the duration of the participant's trial participation

Laboratory Inclusion Values

Immunology/Virology

14. **HIV-1 infection**, with reactive HIV-1 antibody and any Multispot or Geenius HIV-1/HIV-2 results, documented by the HVTN 703/HPTN 081 HIV diagnostic algorithm
15. **Plasma HIV-1 RNA $\geq 1,000$ copies/mL** by any assay, prior to initiating ART
16. **CD4+ T-cell count ≥ 450 cells/mm³** obtained within 90 days prior to enrollment
17. One **plasma HIV-1 RNA below the lower limit of quantitation (LLOQ)** collected at each of the following:
 - at screening, within 90 days prior to enrollment; and
 - greater than 9 months prior to the screening HIV-1 RNA

Note: Sites must have results from locally available assays that are approved as standard-of-care by their regional governing bodies.

Hematology

18. **Hemoglobin (Hgb) ≥ 10.0 g/dL**
19. **Absolute neutrophil count (ANC) ≥ 750 cells/mm³**
20. **Platelets $\geq 100,000$ cells/mm³**

Chemistry

21. **Alanine aminotransferase (ALT)** < 2.5 times the institutional upper limit of normal and **direct bilirubin** within the institutional range of normal
22. **eGFR** > 60 mL/min/1.73m²

Reproductive Status

23. **Volunteers capable of becoming pregnant:** negative serum or urine beta human chorionic gonadotropin (β -HCG) pregnancy test performed at the screening visit and prior to enrollment. Persons who are NOT capable of becoming pregnant due to having reached menopause (no menses for 1 year) or having undergone total hysterectomy or bilateral oophorectomy or tubal ligation (verified by medical records) are not required to undergo pregnancy testing.
24. **Reproductive status:** A volunteer who is capable of becoming pregnant must agree to consistently use effective contraception (ie, IUD or hormonal; see [Appendix B](#) and HVTN 805/HPTN 093 SSP) for sexual activity that could lead to pregnancy from at least 21 days prior to enrollment through confirmation of viral suppression following ART re-initiation.
25. **Volunteers capable of becoming pregnant must also agree not to seek pregnancy through alternative methods**, such as artificial insemination or in vitro fertilization, until after confirmation of viral suppression following ART re-initiation.

5.2 Exclusion criteria

1. Any **plasma HIV-1 RNA** $\geq$ **LLOQ** (LLOQ: 75, 50, 40, or 20 copies/mL) within 12 months prior to enrollment
 - NOTE: Two “blips” (ie, plasma HIV-1 RNA > LLOQ) < 400 copies/mL are allowed if preceded and followed by values < LLOQ and if the blips occur more than 6 months prior to enrollment.

Note: Sites must have results from locally available assays that are approved as standard-of-care by their regional governing bodies.

2. **History of AIDS-defining illnesses** or US Centers for Disease Control (CDC) **Category C events** per the current list on the CDC website (see HVTN 805/HPTN 093 SSP)
3. **Autoimmune disease**, including Type I diabetes mellitus (Not excluded from participation: Volunteer with mild, stable and uncomplicated autoimmune disease that does not require consistent immunosuppressive medication and that, in the judgment of the site investigator, is likely not subject to exacerbation and likely not to complicate AE assessments).

4. **Immunosuppressive medications** received within 6 months before enrollment (Not exclusionary: [1] corticosteroid nasal spray; [2] inhaled corticosteroids; [3] topical corticosteroids for mild, uncomplicated dermatologic condition; or [4] a single course of oral/parenteral prednisone or equivalent at doses ≤ 60 mg/day and length of therapy < 11 days with completion at least 30 days prior to enrollment)
5. **Blood products** received within 120 days before planned ART interruption
6. **Investigational research agents**, other than experimental vaccine(s) (See Exclusion Criterion #7), received within 30 days before planned ART interruption
7. **HIV or non-HIV experimental vaccine(s) received within the last 1 year.** Exceptions may be made by the HVTN 805/HPTN 093 Protocol Safety Review Team (PSRT) for vaccines that have subsequently undergone licensure by the FDA or by the national regulatory authority where the volunteer is enrolling. For volunteers who have received control/placebo in an experimental vaccine trial, the HVTN 805/HPTN 093 PSRT will determine eligibility on a case-by-case basis. For volunteers who have received an experimental vaccine(s) greater than 1 year ago, eligibility for enrollment will be determined by the HVTN 805/HPTN 093 PSRT on a case-by-case basis.
8. **Licensed live attenuated vaccines** received within 30 days before planned ART interruption (eg, measles, mumps, and rubella [MMR]; oral polio vaccine [OPV]; varicella; yellow fever; live attenuated influenza vaccine)
9. **Licensed vaccines that are not live attenuated vaccines** received within 14 days before planned ART interruption (eg, tetanus, pneumococcal, hepatitis A or B, influenza)
10. **Significant unstable cardiac or cerebrovascular disease** (eg, angina, congestive heart failure [CHF], recent cerebrovascular accident [CVA], or myocardial infarction [MI])
11. **Positive Hepatitis B surface antigen (HBsAg) or positive HCV RNA** (Not exclusionary: positive HCV Ab with negative HCV RNA)
12. **Pregnant or breastfeeding**
13. **Clinically significant medical condition**, physical examination findings, clinically significant abnormal laboratory results, or past medical history with clinically significant implications for current health. A clinically significant condition or process includes but is not limited to:
 - A process that would affect the immune response;
 - A process that would require medication that affects the immune response;

- Any contraindication to repeated blood draws, including inability to establish venous access;
 - A condition that requires active medical intervention or monitoring to avert grave danger to the volunteer's health or well-being during the study period; or
 - Any condition specifically mentioned among the exclusion criteria.
14. **Any medical, psychiatric, occupational, or other condition** that, in the judgment of the investigator, would interfere with, or serve as a contraindication to, protocol adherence, assessment of safety, or a volunteer's ability to give informed consent
 15. **Any medical, psychiatric, occupational, or other condition** that, in the judgment of the investigator, could be exacerbated by events associated with protocol participation, which include: ATI, low-level viremia, subsequent viral rebound, and ART re-initiation
 16. **HIV dementia or other neurologic disease** that, in the judgment of the investigator, would be a contraindication to study participation
 17. **Psychiatric condition that precludes compliance with the protocol.** Specifically excluded are persons with psychoses within the past 3 years, ongoing risk for suicide, or history of suicide attempt or gesture within the past 3 years.
 18. **Malignancy** (Not excluded from participation: Volunteer who has had malignancy excised surgically and who, in the investigator's judgment, has a reasonable assurance of sustained cure, or who is unlikely to experience recurrence of malignancy during the period of the study)
 19. **Current untreated or incompletely treated active tuberculosis disease or current latent tuberculosis infection** (Not excluded from participation: Volunteer who has latent tuberculosis infection and is undergoing treatment, with at least one month of treatment completed)
 20. **Untreated or incompletely treated syphilis, gonorrhea, or chlamydia infection**

5.3 Criteria for participant early termination

- Pregnancy or breastfeeding (see Section [6.7.1](#))
- Declared intention to become pregnant or begin breastfeeding before confirmation of viral suppression following ART re-initiation
- Request by the participant to withdraw

- Request of the primary HIV care provider
- Participant starts ART switch but is not virally suppressed after approximately 12 weeks (see Section 6.1.3).
- Participant is judged by the investigator to be at significant risk of failing to comply with the provisions of the protocol so as to cause to harm to self or seriously interfere with the validity of the study results.
- Participant relocates and remote follow-up or transfer to another CRS is not possible. (Note: Remote follow-up is allowable only after confirmation of viral suppression after ART re-initiation; see Ab Manual of Operations (MOP) and HVTN 805/HPTN 093 SSP for further detail.)
- CRS determines that the participant is lost to follow-up.
- Investigator decides, in consultation with Protocol Team leadership, to terminate participation (eg, if participant exhibits inappropriate behavior toward clinic staff).
- Any condition where termination from the study is required by applicable regulations

5.4 Study termination

This study may be terminated early by the determination of the HVTN 805/HPTN 093 PSRT, a pertinent national regulatory authority, NIH, or the Office for Human Research Protections (OHRP). In addition, the conduct of this study at an individual Network CRS may be terminated by the determination of the IRB/EC and any applicable RE.

6 Study procedures

For all study visits, unless otherwise specified, participants will come to their assigned CRS.

6.1 Schedule 1: Monitoring ATI

6.1.1 Screening

Screening may occur over the course of several contacts/visits. All inclusion and exclusion criteria must be assessed within 8 weeks (56 days) prior to enrollment, unless otherwise specified in the eligibility criteria (Section 5).

After signing informed consent, volunteers will undergo the following procedures, as shown in [Appendix E](#) and [Appendix H](#):

- Assessment of Understanding
- Medical history, including:
 - current psychiatric status and history
 - assessment of routine clinically-indicated vaccinations (note: must be completed prior to ATI initiation (see Section 5.2))
- Complete physical examination
- Assessment of concomitant medications
- HIV transmission risk behavior assessment and counseling
- Contraception status assessment (for participants who are capable of becoming pregnant and who are sexually active in a way that could lead to pregnancy; see 6.6 and [Appendix B](#))
- Decision aid
- Blood collection for:
 - HIV PCR viral load
 - CD4+ and CD8+ T-cell counts
 - HBsAg and hepatitis C serology
 - QuantiFERON TB test

- o Hgb, ANC, platelets
- o ALT, direct bilirubin, eGFR
- o Syphilis testing
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause (no menses for 1 year), or having undergone hysterectomy or bilateral oophorectomy or tubal ligation [verified by medical records], are not required to undergo pregnancy testing)
- Gonorrhea/chlamydia testing by urine or cervical/vaginal swab
- Trichomonas testing by cervical/vaginal swab
- Tuberculin skin test (TST; only if QuantiFERON TB test not available)

6.1.2 Enrollment

- Participants undergoing an “ART switch” will be considered enrolled on the first day of the new ART medication.
- Participants who do not undergo an “ART switch” will be considered enrolled at the initiation of ATI (Day 0).

6.1.3 ART Switch

ART switch

Participants taking NNRTI-based ART regimens are required to switch to a PI- or INSTI-based regimen at least 4 weeks before beginning ATI and must demonstrate viral suppression (ie, < LLOQ) on the new regimen before beginning ATI.

At the clinic visit initiating the ART switch, the participant will undergo the following procedures (see [Appendix E](#) and [Appendix H](#)):

- Targeted (ie, symptom directed) physical examination, including weight and vital signs
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- HIV transmission risk behavior assessment and counseling

- Contraception status assessment (for participants who are capable of becoming pregnant and who are sexually active in a way that could lead to pregnancy; see [Section 6.6](#) and [Appendix B](#))
- Decision-making assessment
- Psychosocial assessment
- Blood collection for:
 - o HIV PCR viral load
 - o CD4+ and CD8+ T-cell counts
 - o Hgb, ANC, platelets
 - o ALT, direct bilirubin, eGFR
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause (no menses for 1 year), or having undergone hysterectomy or bilateral oophorectomy or tubal ligation [verified by medical records], are not required to undergo pregnancy testing)

Post-ART switch contact

The site will make arrangements to contact the participant approximately 2 weeks after initiation of the new ART regimen to assess tolerability of the ART switch and willingness to continue. This contact may be conducted over the phone or in person; see HVTN 805/HPTN 093 SSP for details.

ATI qualification visit

Four weeks or more after initiation of the new ART regimen, participants undergoing an ART switch will return for a clinic visit during which they will undergo the following procedures (see [Appendix E](#) and [Appendix H](#)):

- Targeted (ie, symptom directed) physical examination, including weight and vital signs
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- HIV transmission risk behavior assessment and counseling
- Contraception status assessment (for participants who are sexually active in a way that could lead to pregnancy; see [Appendix B](#))

- Blood collection for:
 - o HIV PCR viral load
 - o CD4+ and CD8+ T-cell counts
 - o Hgb, ANC, platelets
 - o ALT, direct bilirubin, eGFR
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause (no menses for 1 year), or having undergone hysterectomy or bilateral oophorectomy or tubal ligation [verified by medical records], are not required to undergo pregnancy testing)

The procedures for the ATI qualification visit should be repeated as appropriate until viral suppression (ie, < LLOQ) is demonstrated. In general, ATI should be initiated approximately 2 weeks after the ATI qualification visit at which samples are collected that demonstrate viral suppression on the new ART regimen.

Any participant who is not virally suppressed at the ATI qualification visit will be retested on a weekly basis until virally suppressed. If not virally suppressed by approximately 12 weeks from the start of the ART switch, the participant should be referred to re-initiate their previous NNRTI-containing regimen, will not proceed to the ATI, and will be terminated from the study.

6.1.4 ATI

For all participants, initiation of ATI is defined as Day 0.

While on Schedule 1, participants will undergo the following procedures, as specified in [Appendix E](#) and [Appendix H](#):

- Complete physical examination OR Targeted (ie, symptom directed) physical examination, including weight and vital signs
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- ART re-initiation assessment
- HIV transmission risk behavior assessment and counseling
- Contraception status assessment (for participants who are capable of becoming pregnant and who are sexually active in a way that could lead to pregnancy; see Section [6.6](#) and [Appendix B](#))

- Decision-making assessment
- Psychosocial assessment
- Social impact assessment
- Social impact assessment questionnaire
- Blood collection per [Appendix E](#) for:
 - o HIV PCR viral load
 - o CD4+ and CD8+ T-cell counts
 - o Hgb, ANC, platelets
 - o ALT, direct bilirubin, eGFR
 - o Syphilis testing
 - o ARV detection
 - o Intracellular cytokine staining (ICS)
 - o Immune cell phenotyping
 - o Neutralizing antibodies (nAb)
 - o Fragment crystallizable receptor (FcR)-mediated effector functions
 - o HIV reservoir assessment
 - o Blood hormone levels (estradiol and progesterone)
 - o Serum, plasma, and PBMC storage
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause (no menses for 1 year), or having undergone hysterectomy or bilateral oophorectomy or tubal ligation [verified by medical records], are not required to undergo pregnancy testing)
- Gonorrhea/chlamydia testing by urine or cervical/vaginal swab
- Trichomonas testing by cervical/vaginal swab

6.2 Schedule 2: Monitoring ATI with viremia

As soon as participants demonstrate viral load ≥ 200 copies/mL, they will transition to Schedule 2, during which they will continue ATI while viremia is monitored. At Schedule 2 timepoints specified in [Appendix F](#) and [Appendix I](#), participants will undergo the following procedures:

- Complete physical examination OR targeted (ie, symptom directed) physical examination, including weight and vital signs
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- ART re-initiation assessment
- HIV transmission risk behavior assessment and counseling
- Contraception status assessment (for participants who are capable of becoming pregnant and who are sexually active in a way that could lead to pregnancy; see Section 6.6 and [Appendix B](#)).
- Decision-making assessment
- Psychosocial assessment
- Social impact assessment
- Social impact assessment questionnaire
- Blood collection for:
 - o HIV PCR viral load
 - o CD4+ and CD8+ T-cell counts
 - o Hgb, ANC, platelets
 - o ALT, direct bilirubin, eGFR
 - o Syphilis testing
 - o ARV detection
 - o ICS
 - o Immune cell phenotyping

- o Neutralizing antibodies (nAb)
- o FcR-mediated effector functions
- o Blood hormone levels (estradiol and progesterone)
- o Serum, plasma, and PBMC storage
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause [no menses for 1 year] or having undergone hysterectomy or bilateral oophorectomy [verified by medical records], are not required to undergo pregnancy testing).
- Gonorrhea/chlamydia testing by urine or cervical/vaginal swab
- Trichomonas testing by cervical/vaginal swab

6.3 Schedule 3: Follow-up on ART

After ART re-initiation criteria are met (see Sections [3.3.2](#) and [3.3.3](#)), participants will transition to Schedule 3 once ART is re-initiated (see [Appendix G](#) and [Appendix J](#)). In Schedule 3, if virologic resuppression has not been achieved by Week 12, participants may continue with biweekly visits until viral resuppression is achieved. These additional visits will be treated as interim visits (see HVTN 805/HPTN 093 SSP).

At Schedule 3 timepoints specified in [Appendix G](#) and [Appendix J](#), participants will undergo the following procedures:

- Complete physical examination OR targeted (ie, symptom directed) physical examination, including weight and vital signs
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- HIV transmission risk behavior assessment and counseling
- Contraception status assessment (for participants who are capable of becoming pregnant and who are sexually active in a way that could lead to pregnancy; see Section [6.6](#) and [Appendix B](#)).
- Decision-making assessment
- Psychosocial assessment

- Social impact assessment
- Social impact assessment questionnaire
- Blood collection for:
 - o HIV PCR viral load
 - o CD4+ and CD8+ T-cell counts
 - o Hgb, ANC, platelets
 - o ALT, direct bilirubin, eGFR
 - o Syphilis testing
 - o HIV genotypic antiretroviral resistance
 - o Viral isolation and sequencing
 - o ICS
 - o Immune cell phenotyping
 - o Neutralizing antibodies (nAb)
 - o FcR-mediated effector functions
 - o HIV reservoir assessment
 - o Serum, plasma, and PBMC storage
- Urine pregnancy test (persons not of reproductive potential due to having reached menopause [no menses for 1 year] or having undergone hysterectomy or bilateral oophorectomy [verified by medical records], are not required to undergo pregnancy testing).
- Gonorrhea/chlamydia testing by urine or cervical/vaginal swab
- Trichomonas testing by cervical/vaginal swab

6.4 Interim visits

Confirmatory VL samples and confirmatory CD4+ T cell samples may be drawn at interim visits (see Sections [3.3.1](#), [3.3.2](#), and [3.3.3](#)). At interim visits for this purpose, the following clinical procedures will be performed:

- Targeted physical exam
- Assessment of concomitant medications
- Intercurrent illness/adverse experiences (AEs)
- ART re-initiation assessment
- HIV transmission risk behavior assessment and counseling
- Social impact assessment

Additional procedures may be performed at the discretion of the clinician.

For additional information on interim visits, including interim visits for other reasons, see the HVTN 805/HPTN 093 SSP.

6.5 Early termination visit

At an early termination visit for a participant on Schedule 1 or 2, CRS staff should consider performing procedures specified for Extended Visit Type A (see [Appendix E](#), [Appendix F](#), [Appendix H](#), and [Appendix I](#)). Such participants will be urged to re-initiate ART under the care of their primary HIV care provider (see HVTN 805/HPTN 093 SSP for additional details).

At an early termination visit for a participant on Schedule 3, CRS staff should consider performing laboratory procedures specified for Week 40 (Visit 91) and clinic procedures specified for Week 52 (Visit 92) (see [Appendix G](#) and [Appendix J](#); see HVTN 805/HPTN 093 SSP for additional details).

6.6 Contraception status

Contraception status is assessed and documented at every scheduled clinic visit from screening through confirmation of viral resuppression following ART re-initiation for a participant who is capable of becoming pregnant and who is sexually active in a way that could cause that participant to become pregnant. Prior to enrollment and until viral resuppression is confirmed following ART re-initiation, staff will ask participants to verbally confirm their use of adequate contraceptive methods (see [Appendix B](#)). A participant who is sexually active in a way that could cause that participant to become pregnant should be reminded at all scheduled clinic visits of the importance of using effective contraception (ie, IUD or hormonal) in addition to barrier protection (ie, condoms) to prevent HIV transmission to sexual partner(s) and should be referred to specific counseling, information, and advice as needed. This reminder should be documented in the participant's study record.

Infertility—including having reached menopause (no menses for 1 year) or having undergone hysterectomy, bilateral oophorectomy, or tubal ligation—must be documented in the participant’s study record.

6.7 Specific clinical management considerations

6.7.1 Pregnancy or breastfeeding

Volunteers who are pregnant or breastfeeding are not eligible to participate in this study. Participants who become pregnant or are breastfeeding during the ART switch will not be allowed to interrupt ART and will be terminated from the study immediately. Participants who become pregnant or are breastfeeding during the ATI phase will re-initiate ART as soon as possible. Post-ATI follow-up will continue per Schedule 3 for up to 24 weeks after viral re-suppression is confirmed following ART re-initiation.

Participants who become pregnant or begin breastfeeding while on Schedule 3 (ie, after ART re-initiation) will continue per Schedule 3 for up to 24 weeks after viral re-suppression is confirmed.

If a participant completes the study or chooses to discontinue from the study before the pregnancy ends, then CRS staff should request permission to contact the participant regarding pregnancy outcomes at the end of the pregnancy. If obtained, pregnancy outcomes will be recorded on a CRF at the end of the pregnancy. All pregnancies must be reported to the Antiretroviral Pregnancy Registry.

6.7.2 Acute retroviral syndrome (ARS)

ARS is a rare diagnosis to be made by a CRS clinical IoR or designee or a primary HIV care provider. The HVTN 805/HPTN 093 PSRT should be consulted on possible ARS diagnosis. Participants with this diagnosis will re-initiate ART as soon as possible.

Signs and symptoms that may support a diagnosis of ARS include, but are not limited to, unintentional weight loss > 5% of pre-ATI body weight, otherwise unexplained persistent fever [$> 38^{\circ}\text{C}$], persistent night sweats, persistent diarrhea, oral candidiasis, and generalized lymphadenopathy.

7 Study products

There are no study products. Drugs for ART and PrEP will not be provided by the study or paid for using sponsor funds. Access to external funding sources for PrEP and ART provision is available. Procedures for accessing external funding sources for PrEP and ART provision are detailed in the HVTN 805/HPTN 093 SSP.

8 Laboratory

8.1 CRS laboratory procedures

The HVTN 805/HPTN 093 Site Lab Instructions and HVTN 805/HPTN 093 SSP provide further guidelines for operational issues concerning the clinical and processing laboratories. These documents include guidelines for general specimen collection, special considerations for phlebotomy, specimen labeling, whole blood processing, HIV screening/diagnostic testing, and general screening and safety testing.

Tube types for blood collection are specified in [Appendix E](#), [Appendix F](#), and [Appendix G](#). For tests performed locally, the local lab may assign appropriate tube types.

In specific situations, the blood collection tubes may be redirected to another laboratory or may require study-specific processing techniques. In these cases, laboratory special instructions will be posted on the protocol-specific section of the HVTN website.

Of note, all assays described below with the exception of HIV antiretroviral resistance testing are performed as research assays and are not approved for use in medical care. Results from these assays are not made available to participants or medical professionals to guide treatment decisions.

8.2 Assay timepoints

Endpoint assays for immunologic and virologic responses will be performed on a subset of participants to be determined by the outcome of primary objectives and endpoints.

8.3 Endpoint assays: cellular

8.3.1 Intracellular cytokine staining (ICS) assay

Flow cytometry will be used to examine HIV-specific CD4⁺ and CD8⁺ T-cell responses following stimulation of PBMCs with synthetic HIV peptides that span the HIV-1 viral proteins. ICS parameters will include cytokines such as interferon gamma (IFN- γ), interleukin (IL)-2, and TNF- α , and may include other cytokines (such as cytokines relevant to Th2 and Th17 responses) to identify T cells of specific functionality. Data will be reported as percentages of CD4⁺ and CD8⁺ T cells responding to a specific peptide pool. Additional cell surface markers, cytokines, or functional markers may also be analyzed.

8.4 Phenotyping of cell populations

Phenotyping of dendritic cells, monocytes, NK cells, B cells, T cells, or other leukocytes for lineage, maturation, and activation markers by flow cytometry may be performed on PBMCs. Data will be reported as percent of cells positive for each marker at the various timepoints.

8.5 Endpoint assays: humoral

8.5.1 Neutralizing antibody assay

HIV-1–specific nAb assays will be performed on serum samples from study participants. The TZM-bl assay will test neutralization of the autologous transmitted/founder virus and a panel of heterologous tier 2 viruses (118, 119). Autologous neutralization escape may also be assessed at the discretion of the HVTN Laboratory Center, contingent on the results of the nAb assays.

8.5.2 Antibody-dependent cellular cytotoxicity (ADCC)

ADCC activity may be assessed using serum samples from study participants. For the Luciferase-based cytotoxicity assay, participant sera are incubated with infectious molecular clone (IMC)-infected cells and percent killing is measured by evaluating the reduction in the luminescence signal after incubation of infected-target and effector cells in presence of serum Ab.

8.5.3 Antibody-dependent cellular phagocytosis (ADCP)

To assess the ability of antibodies to engage cellular FcR for potential antiviral function, ADCP may be measured using serum samples from study participants. ADCP is measured by assessing the ability of antibodies to mediate monocyte phagocytosis of HIV-1 antigen coated fluorescent beads by flow cytometry (120, 121). An array of antigens or viruses may also be analyzed at the discretion of the HVTN Laboratory Center, which may be contingent on the results of the primary antigens or viruses.

8.5.4 Virion capture

The ability of purified IgG from participant serum to mediate infectious virion capture will be measured by infectious virion capture assay (IVCA). The test uses a Protein G column-based capture of Ig-virion immune complexes.

8.6 HIV viral reservoir

Changes in the HIV reservoir will be quantitated in CD4⁺ T cells purified from PBMCs at designated timepoints, using testing methods such as the Intact Provir

DNA Assay (IPDA), *Tat/rev* Induced Limiting Dilution Assay (TILDA), assays detecting replication-competent virus-bearing cells (eg, virus outgrowth assays), and/or measures of total proviral DNA. Cell-associated HIV-RNA may be quantitated as a measure of the transcriptionally active reservoir.

8.7 Host genomics

HLA and CCR5 genotyping may be performed on samples from this study or HVTN 703/HPTN 081 using cryopreserved PBMC. Other genes, including those associated with immune responses (eg, immunoglobulin or T-cell receptor genes) or HIV-1 disease progression may also be assessed.

8.8 HIV antiretroviral resistance testing

HIV antiretroviral resistance testing will be performed by RNA sequencing for resistance mutations in participants whose plasma viremia shows limited suppression after ART re-initiation.

8.9 ARV detection

Blood specimens will be collected during treatment interruption for ARV detection and drug level testing. ARV drug levels may be assessed at the discretion of the protocol team for participants who have not met virologic and non-virologic criteria for transition from monitoring ATI (Schedule 1) to monitoring ATI with viremia (Schedule 2) or for ART re-initiation (from Schedule 2 to Schedule 3). Assay methods may include HPLC-mass spectrometry, reverse-phase HPLC coupled with tandem mass spectrometry, or another similar method.

8.10 Biohazard containment

As the transmission of HIV and other blood-borne pathogens can occur through contact with contaminated needles, blood, and blood products, appropriate precautions will be employed by all personnel in the drawing of blood and shipping and handling of all specimens for this study, as currently recommended by the CDC and the NIH or other applicable agencies.

All dangerous goods materials, including Biological Substances, Category A or Category B, must be transported according to instructions detailed in the International Air Transport Association Dangerous Goods Regulations.

9 Research use of stored human samples and/or data

9.1 Specimen storage and other use of specimens

The HVTN and HPTN store specimens from all study participants indefinitely, unless a participant requests that specimens be destroyed or if required by IRB/EC, or RA.

Other use of specimens is defined as studies not covered by the protocol or the informed consent form for the main study (see [Appendix A](#) and [Appendix C](#)).

This research may relate to HIV, vaccines, antibodies, the immune system, and other diseases. This could include genetic testing and, potentially, genome-wide studies. This research is done only to the extent authorized in each study site's informed consent form, or as otherwise authorized under applicable law. Other research on specimens ("other use") will occur only after review and approval by the Networks, the IRB/EC of the researcher requesting the specimens, and the IRBs/ECs/REs of the CRSs if required.

As part of consenting for the study, participants document their initial decision to allow or not allow their specimens to be used in other research, and they may change their decision at any time. The participant's initial decision about other use of their specimens, and any later change to that decision, is recorded by their CRS in a Web-based tool that documents their current decisions for other use of their specimens. The Networks will only allow other research to be done on specimens from participants who allow such use.

CRSs must notify HVTN Regulatory Affairs if institutional or local governmental requirements pose a conflict with or impose restrictions on specimen storage or other use of specimens.

9.2 Exploratory studies

Samples may be used for other testing and research related to furthering the understanding of HIV, immunology, antibodies, or vaccines. In addition, cryopreserved samples may be used to perform additional assays to support standardization and validation of existing or newly developed methods.

10 Potential risks and benefits

10.1 Potential risks

10.1.1 Risks of ATI

The risks from a closely monitored ATI are minimal in this study population. There is a theoretical risk that such an interruption could lead to the development of HIV drug resistance (122). This may be a particular concern for individuals taking NNRTIs. However, this potential risk is substantially mitigated by the procedures described in Section 6.1.1. Given the study population as restricted by the eligibility criteria, the frequency of immunological and virologic monitoring, and the criteria for restarting ART, it is extremely unlikely that such an ART interruption will lead to the development of any opportunistic infections or AIDS-defining conditions.

Rarely, viremic rebound may be associated with ARS (see Section 6.7.2).

During the ATI phase, participants may transmit HIV infection if they do not adhere to safer sex practices.

10.1.2 Phlebotomy

Drawing blood may be associated with discomfort, bruising, local hematoma formation and, on rare occasions, infections, lightheadedness, and fainting.

The amount of blood drawn for research purposes will be within regulatory limits (see 21 CFR 630.15).

10.2 Potential benefits

Study participants should expect no direct health benefits from study participation. Knowledge gained in this study may aid in development of better interventions against HIV.

11 Monitoring of safety

11.1 Safety monitoring and oversight

11.1.1 HVTN 805/HPTN 093 PSRT

The HVTN 805/HPTN 093 PSRT is composed of the following members:

- DAIDS medical officer representatives
- Protocol chairs
- Protocol Team leaders
- Core medical monitor
- Clinical safety specialist
- Regional medical liaison

The Protocol Team clinic coordinator, clinical data manager, clinical trial manager, clinical research manager, and others may also be included in HVTN 805/HPTN 093 PSRT meetings.

The clinician members of HVTN 805/HPTN 093 PSRT are responsible for decisions related to participant safety.

The PSRT will review laboratory parameters (HIV-RNA, CD4+ T-cell count), adverse events, and a summary of the distribution of time to meeting ART re-initiation criteria twice a month. Once the final participant achieves viral resuppression in Schedule 3, the PSRT may consider holding less frequent reviews (ie, monthly).

The PSRT will also review the distribution of time to meeting ART re-initiation criteria in the first 20 HVTN 805/HPTN 093 participants and assess whether the distribution is comparable to that observed in an NIH intramural therapeutic HIV vaccine study that enrolled HIV-infected participants who initiated ART early with similar ART re-initiation criteria to those proposed here. Based on the NIH intramural therapeutic HIV vaccine study results, we expect that the median time to meeting ART re-initiation criteria in HVTN 805/HPTN 093 will be at ATI week 15 and that the 75th percentile for time to meeting ART re-initiation criteria will be around week 12-18. If the spread in the distribution time to meeting ART re-initiation criteria is less than expected, then the HVTN 805/HPTN 093 PSRT, in consultation with the NIAID Data and Safety Monitoring Board (DSMB), will consider strategies for modifying the study (eg, to alter the eligibility criteria) with the aim to further enrich for potential controllers.

The HVTN 805/HPTN 093 PSRT will also monitor specifically for failure to re-suppress or to demonstrate a 2-log viral load reduction within 12 weeks after ART re-initiation. Any such failure and/or any AE $\geq$ Grade 4 deemed related to the study procedures will be reviewed by the HVTN 805/HPTN 093 PSRT promptly and, if indicated, by the NIAID DSMB to evaluate whether and how the study should be modified.

11.1.2 NIAID DSMB

The NIAID DSMB assesses the study conduct during the trial and may give advice to the HVTN 805/HPTN 093 Protocol Team leadership.

Approximately 6 months after enrollment of the first participant or after the 10th participant has completed 12 weeks of ATI (whichever comes first), an interim review of the study will occur. The DSMB will review accrual; retention; AE summaries, including all reported AEs $\geq$ Grade 3 and all STIs; summaries of the time to meeting ART re-initiation criteria; and longitudinal summaries of HIV-1 RNA and CD4+ T-cell count. Subsequent DSMB reviews will occur at least annually while participants remain on study. A DSMB may also be convened if a reason is identified by the DAIDS MO, study Co-chairs, or study statisticians in consultation with the HVTN 805/HPTN 093 Protocol Team leadership.

11.1.3 Roles and responsibilities in safety monitoring

The roles and responsibilities of the Statistical and Data Management Center (SDMC) in relation to safety monitoring include:

- Maintaining a central database management system for clinical data;
- Providing reports of clinical data to appropriate groups such as the HVTN 805/HPTN 093 PSRT and NIAID DSMB (see Section [11.1.2](#));

The roles and responsibilities of the HVTN CSS, HVTN RML, or HVTN Core designee in relation to safety monitoring include:

- Daily monitoring of clinical data;
- Querying CRSs for additional information regarding reported clinical data; and
- Providing support to the HVTN 805/HPTN 093 PSRT.

11.2 Safety reporting

11.2.1 Submission of safety forms to SDMC

CRS staff must submit all safety forms (eg, AEs, local lab results, concomitant medications) before the end of the next business day, excluding federal or bank holidays. The forms should not be held in anticipation of additional information at a later date. If additional information is received at a later date, the forms should be updated and resubmitted before the end of the next business day after receiving the new information. For the case of a longer CRS holiday closure, CRS staff must submit the data by the end of the 5th day (local time) after receiving the information even if this day is a holiday.

For example: If the CRS becomes aware of an AE on Thursday (Day 0), the CRS must submit the data by the end of the next business day, on Friday. If there is a longer CRS holiday closure, then this AE must be reported no later than the end of the fifth day, Monday (Day 4). If Monday is a holiday as well, all safety forms still need to be submitted by the end of Monday (Day 4).

11.2.2 Serious adverse event (SAE)

An SAE is an AE that results in one or more of the following outcomes:

- Death
- A life-threatening (ie, an immediate threat to life) event
- Requires in-patient hospitalization or prolongation of an existing hospitalization
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions
- A congenital anomaly/birth defect
- A medically important event (medical and scientific judgment should be exercised in deciding whether expedited reporting is appropriate in such situations, such as important medical events that may not be immediately life threatening or result in death or hospitalization, but which may jeopardize the individual or may require intervention to prevent one of the outcomes listed above.)

11.2.3 AE reporting

An AE is any untoward medical occurrence in a clinical investigation participant, including an abnormal laboratory finding, symptom, or disease temporally

associated with the individual's participation in the research, whether or not related to the research.

The study intervention for which adverse event attribution reporting is required is ATI.

At designated visits, information regarding AEs will be elicited by appropriate questioning and examinations and will be documented in the medical record and in the electronic database.

All AEs are graded according to the Division of AIDS (DAIDS) Table for Grading the Severity of Adult and Pediatric Adverse Events, Corrected Version 2.1, July 2017, available on the DAIDS Regulatory Support Center (RSC) website at <https://rsc.niaid.nih.gov/clinical-research-sites/daids-adverse-event-grading-tables>, except:

- Unintentional Weight Loss is required to be reported as an AE only if it is considered to be potentially deleterious to the participant's health (see HVTN 805/HPTN 093 SSP); and
- Creatinine is required to be reported as an AE only if it is gradable per the increase from local lab upper limit of normal (ULN) parameter. Do not grade elevated creatinine based on the change from the baseline parameter.
- eGFR is required to be reported as an AE only if it is gradable per reported value or if dialysis is needed. Do not grade decreased eGFR based on the change from the baseline parameter (see HVTN 805/HPTN 093 SSP).

All AEs are reported to the SDMC on the appropriate CRF.

Sites are expected to notify HVTN clinical safety staff of any serious safety concern requiring their attention. Telephone numbers and email addresses are found on the protocol home page on the HVTN Members' site (<https://members.hvtn.org/protocols/hvtn805-hptn093>). Concerns requiring immediate attention should be communicated by calling the clinical safety phone. In the case of email notification, clinical safety staff will reply within one business day.

In addition, CRS investigators are required to submit AE information in accordance with IRB/EC and any applicable RE requirements.

12 Ethics/protection of human subjects

12.1 Informed consent process

Informed consent is a process where information is presented to enable persons to voluntarily decide whether or not to participate in a clinical trial. It is an ongoing conversation between the prospective or enrolled research participant and the researchers that begins before consent is given and continues through the decision-making process and until the end of the participant's involvement in the research. Discussions about the research will provide essential information about the study and include: purpose, duration, experimental procedures, alternatives, risks, and benefits. Participants will be given the opportunity to ask questions and to have them answered.

The participant will sign the informed consent document prior to undergoing any research procedures. The participant may withdraw consent at any time throughout the course of the study. A copy of the informed consent document will be given to participants for their records. Documentation of the signing of the consent form will be retained in the participant's medical record. The rights and welfare of participants will be protected by emphasizing to them that the quality of their medical care will not be adversely affected if they decline to participate in this study. These rights and responsibilities are further elaborated upon in the Participants Bill of Rights and Responsibilities developed by the HVTN and HPTN for their jointly conducted trials.

The HVTN 805/HPTN 093 SICF is modeled on previously-approved treatment interruption consent forms and, specifically, emphasizes the cautionary points noted above (see Section 2.6) with respect to the uncertainties, risks and lack of anticipated individual benefits; the fact that treatment interruptions are not recommended in clinical care; and the fact that ART guidelines recommend lifelong treatment. While the SICF is a crucial starting point, the quality of informed consent also relies on the process and other tools employed to supplement that document. With respect to these supplementary measures the team is building upon each Network's longstanding efforts to build rigorous informed consent into trial implementation (123). These efforts begin with the community engagement described above (see Section 2.7) and go on to include: drafting of the SICF by and with local community representatives who reflect the target population and the sites that will be conducting the study; review of translated materials by local staff and community representatives who are native speakers of each language to help ensure accuracy and appropriateness; the provision of a Frequently Asked Questions information sheet; the development and use of visual aids to convey some concepts, such as blood draw volumes; the development of materials and processes to engage sexual partners, if participants and their partners so choose; the use of an Assessment of Understanding and re-review of any misunderstandings as part of the initial and ongoing informed

consent process; and ongoing community engagement throughout the trial, including local education and engagement activities, CAB meetings and study-specific stakeholder meetings, among other forums.

For HVTN 805/HPTN 093, the team will develop additional tools to facilitate optimal informed consent and decision-making. Specifically, the team will develop decision aids, such as those found to improve the quality of shared decision making and consent in preference-sensitive healthcare decisions (124, 125) and will partner with participant advocates—nonteam members, ideally peers, familiar with the AMP Study, treatment interruption, and the HVTN 805/HPTN 093 study—who could work with the AMP participant to further facilitate optimal decision-making regarding potential HVTN 805/HPTN 093 participation. The latter model has been used, for example, in the setting of organ transplantation between HIV-positive donors and recipients (126, 127). Clinic staff will also be in consultation with each participant's primary HIV care provider, as needed, to ensure their understanding and ability to support the participants. Any such materials developed for participants will be submitted for review and approval to the IRB/EC prior to using with participants.

12.2 Participant confidentiality

All records will be kept confidential to the extent provided by federal, state, and local law. Study monitors and other authorized representatives of the Sponsor may inspect all documents and records required to be retained by the investigator, including but not limited to medical records. Records will be kept locked and all computer entry and networking programs will be done with coded numbers only. Clinical information will not be released without written permission of the participant, except as necessary for monitoring by IRB, NIH/NIAID/DAIDS or the sponsor's designee, OHRP, or another regulatory authority.

13 Protocol conduct

This protocol and all actions and activities connected with it will be conducted in compliance with Good Clinical Practice (GCP) (ICHe6), the HVTN and HPTN network-specific *Manuals of Operations*, and DAIDS Clinical Research Policies and Standard Procedures Documents, including procedures for the following:

- Protocol registration, activation, and implementation;
- Informed consent, screening, and enrollment;
- Study participant reimbursement;
- Clinical and safety assessments;
- Safety monitoring and reporting;
- Data collection, documentation, transfer, and storage;
- Participant confidentiality;
- Study follow-up and close-out;
- Quality control;
- Protocol monitoring and compliance;
- Advocacy and assistance to participants regarding negative social impacts associated with the trial;
- Risk reduction counseling;
- Specimen collection, processing, and analysis;
- Exploratory and ancillary studies and sub-studies, and
- Destruction of specimens.

Any policies or procedures that vary from DAIDS, HVTN, or HPTN standards or require additional instructions will be described in the HVTN 805/HPTN 093 SSP.

13.1 Emergency communication with study participants

As in all clinical research, this study may generate a need to reach participants quickly to avoid imminent harm, or to report study findings that may otherwise concern their health or welfare.

When such communication is needed, the CRS will request that its IRB/EC and any applicable RE expedite review of the message. If this review cannot be completed in a timeframe consistent with the urgency of the required communication, the site should contact the participant first, and then notify the IRB/EC and any applicable RE of the matter as soon as possible.

14 Acronyms and abbreviations

ACTG	AIDS Clinical Trials Group
ADCC	antibody-dependent cellular cytotoxicity
ADCP	antibody-dependent cellular phagocytosis
AE	adverse experience
AIDS	acquired immune deficiency syndrome
AMP	antibody mediated prevention
ANC	absolute neutrophil count
ARS	acute retroviral syndrome
ART	antiretroviral therapy
ATI	analytical treatment interruption
AUC	area under the curve
bnAb	broadly neutralizing antibody
CAB	community advisory board
caRNA	cell associated RNA
CBC	complete blood count
CDC	(U.S.) Centers for Disease Control and Prevention
CHI	chronic HIV infection
CRF	case report form
CRS	clinical research site
DAIDS	Division of AIDS
DNA	deoxyribonucleic acid
DSMB	Data and Safety Monitoring Board
EAE	adverse event requiring expedited reporting
EC	ethics committee
eGFR	estimated glomerular filtration rate
EIA	enzyme immunoassay
ELISA	enzyme-linked immunosorbent assay
Fc	fragment crystallizable
Fc γ R	fragment crystallizable gamma receptor
FcR	fragment crystallizable receptor
HAART	highly active antiretroviral therapy
HBsAg	Hepatitis B surface antigen
HBV	Hepatitis B virus
HCV	Hepatitis C virus
Hgb	hemoglobin
HIV	human immunodeficiency virus
HPTN	HIV Prevention Trials Network
HVTN	HIV Vaccine Trials Network
ICS	intracellular cytokine staining
IgG	immunoglobulin G
INSTI	integrase strand transfer inhibitor
IoR	investigator of record
IQR	interquartile range
IRB	institutional review board

LLOQ	lower limit of quantitation
LOC	leadership operations center
mAb	monoclonal antibody
mcL	microliter
nAb	neutralizing antibody
NHP	nonhuman primate
NIAID	National Institute of Allergy and Infectious Diseases
NK	natural killer (cells)
NNRTI	nonnucleoside reverse transcriptase inhibitor
NIH	National Institutes of Health
OHRP	Office of Human Research Protection
PBMC	peripheral blood mononuclear cells
PCR	polymerase chain reaction
PHI	primary HIV infection
PI	protease inhibitor
PK	pharmacokinetic
PrEP	pre-exposure prophylaxis
PSRT	Protocol Safety Review Team
PTC	post-treatment controller
RNA	ribonucleic acid
SAE	serious adverse event
SDMC	statistics and data management center
SHIV	simian/human immunodeficiency virus
SICF	sample informed consent form
SIVIG	specific intravenous immunoglobulin
SMART	Strategies for Management of Antiretroviral Therapy
SSP	study specific procedures
TMLE	targeted minimum loss-based estimation
VL	viral load
VQA	Virology Quality Assurance

15 Protocol version history

Date: April 27, 2020

Protocol version: 1.0

Protocol modification: NA

16 Document references (other than literature citations)

Other documents referred to in this protocol, and containing information relevant to the conduct of this study, include:

- Antiretroviral Pregnancy Registry. Available at <http://www.apregistry.com/>.
- Assessment of Understanding. Accessible through the HVTN protocol-specific website.
- Division of AIDS (DAIDS) Clinical Research Policies and Standard Procedures Documents. Available at <https://www.niaid.nih.gov/research/daids-clinical-research-policies-standard-procedures>
- Division of AIDS Protocol Registration Manual. Available at <https://www.niaid.nih.gov/sites/default/files/prmanual.pdf>
- Division of AIDS Table for Grading the Severity of Adult and Pediatric Adverse Events. Version 2.1, dated March 2017. Available at <https://rsc.niaid.nih.gov/clinical-research-sites/daids-adverse-event-grading-tables>
- The Manual for Expedited Reporting of Adverse Events to DAIDS. Version 2.0, January 2010. Available at <https://rsc.niaid.nih.gov/clinical-research-sites/manual-expedited-reporting-adverse-events-daids>
- HVTN Certificate of Confidentiality. Accessible through the HVTN website.
- HVTN 805/HPTN 093 Study Specific Procedures. Accessible through the HVTN protocol-specific website.
- Ab Manual of Operations. Accessible through the HVTN website.
- HVTN Manual of Operations. Accessible through the HVTN website.
- Dangerous Goods Regulations (updated annually), International Air Transport Association. Available for purchase at <http://www.iata.org/publications/dgr/Pages/index.aspx>
- International Conference for Harmonisation of Technical Requirements for pharmaceuticals for Human Use (ICH) E6 (R2), Integrated Addendum to ICH E6(R1): Guideline for Good Clinical Practice: Section 4.8, Informed Consent of Trial Subjects. Available at https://database.ich.org/sites/default/files/E6_R2_Addendum.pdf

- HVTN 805/HPTN 093 Participants' Bill of Rights and Responsibilities. Accessible through the HVTN website.
- NIH Policy on Reporting Race and Ethnicity Data: Subjects in Clinical Research. Available at <http://grants1.nih.gov/grants/guide/notice-files/NOT-OD-01-053.html>
- Requirements for Source Documentation in DAIDS Funded and/or Sponsored Clinical Trials. Available at <https://www.niaid.nih.gov/labsandresources/resources/daidsclinrsrch/Pages/ClinicalSite.aspx>
- Title 21, Code of Federal Regulations, Part 50. Available at <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=50>
- Title 45, Code of Federal Regulations, Part 46. Available at <http://www.hhs.gov/ohrp/humansubjects/guidance/45cfr46.html>

See Section 17 for literature cited in the background and statistics sections of this protocol.

17 References

1. Gossez M, Martin GE, Pace M, Ramjee G, Premraj A, Kaleebu P, et al. Virological remission after antiretroviral therapy interruption in female African HIV seroconverters. *AIDS*. 2019;33(2):185-97.
2. Treatment Action Group. Community Recommendations for Clinical Research Involving Antiretroviral Treatment Interruptions in Adults Nov 2018 [Available from: http://www.treatmentactiongroup.org/sites/default/files/community_recs_clinical_research_final.pdf.
3. Julg B, Dee L, Ananworanich J, Barouch DH, Bar K, Caskey M, et al. Recommendations for analytical antiretroviral treatment interruptions in HIV research trials-report of a consensus meeting. *Lancet HIV*. 2019;6(4):e259-e68.
4. Chun TW, Fauci AS. HIV reservoirs: pathogenesis and obstacles to viral eradication and cure. *AIDS*. 2012;26(10):1261-8.
5. Chun TW, Stuyver L, Mizell SB, Ehler LA, Mican JA, Baseler M, et al. Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy. *Proc Natl Acad Sci U S A*. 1997;94(24):13193-7.
6. Finzi D, Hermankova M, Pierson T, Carruth LM, Buck C, Chaisson RE, et al. Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science*. 1997;278(5341):1295-300.
7. Wong JK, Hezareh M, Gunthard HF, Havlir DV, Ignacio CC, Spina CA, et al. Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. *Science*. 1997;278(5341):1291-5.
8. Li JZ, Etemad B, Ahmed H, Aga E, Bosch RJ, Mellors JW, et al. The size of the expressed HIV reservoir predicts timing of viral rebound after treatment interruption. *AIDS*. 2016;30(3):343-53.
9. Steingrover R, Pogany K, Fernandez Garcia E, Jurriaans S, Brinkman K, Schuitemaker H, et al. HIV-1 viral rebound dynamics after a single treatment interruption depends on time of initiation of highly active antiretroviral therapy. *AIDS*. 2008;22(13):1583-8.
10. Le CN, Britto P, Brummel SS, Hoffman RM, Li JZ, Flynn PM, et al. Time to viral rebound and safety after antiretroviral treatment interruption in postpartum women compared with men. *AIDS*. 2019;33(14):2149-56.
11. Lodi S, Meyer L, Kelleher AD, Rosinska M, Ghosn J, Sannes M, et al. Immunovirologic control 24 months after interruption of antiretroviral therapy initiated close to HIV seroconversion. *Arch Intern Med*. 2012;172(16):1252-5.

12. Fidler S, Olson AD, Bucher HC, Fox J, Thornhill J, Morrison C, et al. Virological Blips and Predictors of Post Treatment Viral Control After Stopping ART Started in Primary HIV Infection. *J Acquir Immune Defic Syndr*. 2017;74(2):126-33.
13. Goujard C, Girault I, Rouzioux C, Lecuroux C, Deveau C, Chaix ML, et al. HIV-1 control after transient antiretroviral treatment initiated in primary infection: role of patient characteristics and effect of therapy. *Antivir Ther*. 2012;17(6):1001-9.
14. Saez-Cirion A, Bacchus C, Hocqueloux L, Avettand-Fenoel V, Girault I, Lecuroux C, et al. Post-treatment HIV-1 controllers with a long-term virological remission after the interruption of early initiated antiretroviral therapy ANRS VISCONTI Study. *PLoS Pathog*. 2013;9(3):e1003211.
15. Hocqueloux L, Prazuck T, Avettand-Fenoel V, Lafeuillade A, Cardon B, Viard JP, et al. Long-term immunovirologic control following antiretroviral therapy interruption in patients treated at the time of primary HIV-1 infection. *AIDS*. 2010;24(10):1598-601.
16. Stohr W, Fidler S, McClure M, Weber J, Cooper D, Ramjee G, et al. Duration of HIV-1 viral suppression on cessation of antiretroviral therapy in primary infection correlates with time on therapy. *PLoS One*. 2013;8(10):e78287.
17. Martin GE, Gossez M, Williams JP, Stohr W, Meyerowitz J, Leitman EM, et al. Post-treatment control or treated controllers? Viral remission in treated and untreated primary HIV infection. *AIDS*. 2017;31(4):477-84.
18. Namazi G, Fajnzylber JM, Aga E, Bosch RJ, Acosta EP, Sharaf R, et al. The Control of HIV After Antiretroviral Medication Pause (CHAMP) Study: Posttreatment Controllers Identified From 14 Clinical Studies. *J Infect Dis*. 2018;218(12):1954-63.
19. Gianella S, von Wyl V, Fischer M, Niederoest B, Battegay M, Bernasconi E, et al. Effect of early antiretroviral therapy during primary HIV-1 infection on cell-associated HIV-1 DNA and plasma HIV-1 RNA. *Antivir Ther*. 2011;16(4):535-45.
20. Maenza J, Tapia K, Holte S, Stekler JD, Stevens CE, Mullins JI, et al. How often does treatment of primary HIV lead to post-treatment control? *Antivir Ther*. 2015;20(8):855-63.
21. Sneller MC, Justement JS, Gittens KR, Petrone ME, Clarridge KE, Proschan MA, et al. A randomized controlled safety/efficacy trial of therapeutic vaccination in HIV-infected individuals who initiated antiretroviral therapy early in infection. *Sci Transl Med*. 2017;9(419).
22. Martin GE, Frater J. Post-treatment and spontaneous HIV control. *Curr Opin HIV AIDS*. 2018;13(5):402-7.

23. Sharaf R, Lee GQ, Sun X, Etemad B, Aboukhater LM, Hu Z, et al. HIV-1 proviral landscapes distinguish posttreatment controllers from noncontrollers. *J Clin Invest*. 2018;128(9):4074-85.
24. Goulder P, Deeks SG. HIV control: Is getting there the same as staying there? *PLoS Pathog*. 2018;14(11):e1007222.
25. Rouzioux C, Hocqueloux L, Saez-Cirion A. Posttreatment controllers: what do they tell us? *Curr Opin HIV AIDS*. 2015;10(1):29-34.
26. Hurst J, Hoffmann M, Pace M, Williams JP, Thornhill J, Hamlyn E, et al. Immunological biomarkers predict HIV-1 viral rebound after treatment interruption. *Nat Commun*. 2015;6:8495.
27. Gray ES, Taylor N, Wycuff D, Moore PL, Tomaras GD, Wibmer CK, et al. Antibody specificities associated with neutralization breadth in plasma from human immunodeficiency virus type 1 subtype C-infected blood donors. *J Virol*. 2009;83(17):8925-37.
28. Walker LM, Simek MD, Priddy F, Gach JS, Wagner D, Zwick MB, et al. A limited number of antibody specificities mediate broad and potent serum neutralization in selected HIV-1 infected individuals. *PLoS Pathog*. 2010;6(8):e1001028.
29. Stamatatos L, Morris L, Burton DR, Mascola JR. Neutralizing antibodies generated during natural HIV-1 infection: good news for an HIV-1 vaccine? *Nat Med*. 2009;15(8):866-70.
30. Mascola JR, Haynes BF. HIV-1 neutralizing antibodies: understanding nature's pathways. *Immunol Rev*. 2013;254(1):225-44.
31. Kwong PD, Mascola JR. Human antibodies that neutralize HIV-1: identification, structures, and B cell ontogenies. *Immunity*. 2012;37(3):412-25.
32. McCoy LE, Burton DR. Identification and specificity of broadly neutralizing antibodies against HIV. *Immunol Rev*. 2017;275(1):11-20.
33. Burton DR, Hangartner L. Broadly Neutralizing Antibodies to HIV and Their Role in Vaccine Design. *Annu Rev Immunol*. 2016;34:635-59.
34. Zhou T, Georgiev I, Wu X, Yang ZY, Dai K, Finzi A, et al. Structural basis for broad and potent neutralization of HIV-1 by antibody VRC01. *Science*. 2010;329(5993):811-7.
35. Burton DR, Ahmed R, Barouch DH, Butera ST, Crotty S, Godzik A, et al. A Blueprint for HIV Vaccine Discovery. *Cell Host Microbe*. 2012;12(4):396-407.
36. Walker LM, Burton DR. Rational antibody-based HIV-1 vaccine design: current approaches and future directions. *Curr Opin Immunol*. 2010;22(3):358-66.

37. Burton DR, Stanfield RL, Wilson IA. Antibody vs. HIV in a clash of evolutionary titans. *Proc Natl Acad Sci U S A*. 2005;102(42):14943-8.
38. Fauci AS. An HIV Vaccine: Mapping Uncharted Territory. *JAMA*. 2016;316(2):143-4.
39. Haynes BF, Mascola JR. The quest for an antibody-based HIV vaccine. *Immunol Rev*. 2017;275(1):5-10.
40. Halper-Stromberg A, Nussenzweig MC. Towards HIV-1 remission: potential roles for broadly neutralizing antibodies. *J Clin Invest*. 2016;126(2):415-23.
41. Pegu A, Asokan M, Wu L, Wang K, Hataye J, Casazza JP, et al. Activation and lysis of human CD4 cells latently infected with HIV-1. *Nat Commun*. 2015;6:8447.
42. Sloan DD, Lam CY, Irrinki A, Liu L, Tsai A, Pace CS, et al. Targeting HIV Reservoir in Infected CD4 T Cells by Dual-Affinity Re-targeting Molecules (DARTs) that Bind HIV Envelope and Recruit Cytotoxic T Cells. *PLoS Pathog*. 2015;11(11):e1005233.
43. Caskey M, Klein F, Nussenzweig MC. Broadly Neutralizing Antibodies for HIV-1 Prevention or Immunotherapy. *N Engl J Med*. 2016;375(21):2019-21.
44. Margolis DM, Koup RA, Ferrari G. HIV antibodies for treatment of HIV infection. *Immunol Rev*. 2017;275(1):313-23.
45. Ananworanich J, McSteen B, Robb ML. Broadly neutralizing antibody and the HIV reservoir in acute HIV infection: a strategy toward HIV remission? *Curr Opin HIV AIDS*. 2015;10(3):198-206.
46. Ledgerwood JE, Coates EE, Yamshchikov G, Saunders JG, Holman L, Enama ME, et al. Safety, pharmacokinetics and neutralization of the broadly neutralizing HIV-1 human monoclonal antibody VRC01 in healthy adults. *Clin Exp Immunol*. 2015;182(3):289-301.
47. Lynch RM, Boritz E, Coates EE, DeZure A, Madden P, Costner P, et al. Virologic effects of broadly neutralizing antibody VRC01 administration during chronic HIV-1 infection. *Sci Transl Med*. 2015;7(319):319ra206.
48. Bar KJ, Sneller MC, Harrison LJ, Justement JS, Overton ET, Petrone ME, et al. Effect of HIV Antibody VRC01 on Viral Rebound after Treatment Interruption. *N Engl J Med*. 2016;375(21):2037-50.
49. Huang Y, Zhang L, Ledgerwood J, Grunenberg N, Bailer R, Isaacs A, et al. Population pharmacokinetics analysis of VRC01, an HIV-1 broadly neutralizing monoclonal antibody, in healthy adults. *MAbs*. 2017;1-9.
50. Mayer KH, Seaton KE, Huang Y, Grunenberg N, Isaacs A, Allen M, et al. Safety, pharmacokinetics, and immunological activities of multiple intravenous or subcutaneous

doses of an anti-HIV monoclonal antibody, VRC01, administered to HIV-uninfected adults: Results of a phase 1 randomized trial. *PLoS Med.* 2017;14(11):e1002435.

51. Gilbert PB, Juraska M, deCamp AC, Karuna S, Edupuganti S, Mgodini N, et al. Basis and Statistical Design of the Passive HIV-1 Antibody Mediated Prevention (AMP) Test-of-Concept Efficacy Trials. *Stat Commun Infect Dis.* 2017;9(1).
52. Bolton DL, Pegu A, Wang K, McGinnis K, Nason M, Foulds K, et al. Human Immunodeficiency Virus Type 1 Monoclonal Antibodies Suppress Acute Simian-Human Immunodeficiency Virus Viremia and Limit Seeding of Cell-Associated Viral Reservoirs. *J Virol.* 2016;90(3):1321-32.
53. Nishimura Y, Gautam R, Chun T-W, Sadjadpour R, Foulds KE, Shingai M, et al. Early antibody therapy can induce long-lasting immunity to SHIV. *Nature.* 2017;543:559.
54. Scheid JF, Horwitz JA, Bar-On Y, Kreider EF, Lu CL, Lorenzi JC, et al. HIV-1 antibody 3BNC117 suppresses viral rebound in humans during treatment interruption. *Nature.* 2016;535(7613):556-60.
55. Mendoza P, Gruell H, Nogueira L, Pai JA, Butler AL, Millard K, et al. Combination therapy with anti-HIV-1 antibodies maintains viral suppression. *Nature.* 2018;561(7724):479-84.
56. Lu CL, Murakowski DK, Bournazos S, Schoofs T, Sarkar D, Halper-Stromberg A, et al. Enhanced clearance of HIV-1-infected cells by broadly neutralizing antibodies against HIV-1 in vivo. *Science.* 2016;352(6288):1001-4.
57. Barouch DH, Whitney JB, Moldt B, Klein F, Oliveira TY, Liu J, et al. Therapeutic efficacy of potent neutralizing HIV-1-specific monoclonal antibodies in SHIV-infected rhesus monkeys. *Nature.* 2013;503(7475):224-8.
58. Haigwood NL, Montefiori DC, Sutton WF, McClure J, Watson AJ, Voss G, et al. Passive immunotherapy in simian immunodeficiency virus-infected macaques accelerates the development of neutralizing antibodies. *J Virol.* 2004;78(11):5983-95.
59. Schoofs T, Klein F, Braunschweig M, Kreider EF, Feldmann A, Nogueira L, et al. HIV-1 therapy with monoclonal antibody 3BNC117 elicits host immune responses against HIV-1. *Science.* 2016;352(6288):997-1001.
60. Klein F, Nogueira L, Nishimura Y, Phad G, West AP, Jr., Halper-Stromberg A, et al. Enhanced HIV-1 immunotherapy by commonly arising antibodies that target virus escape variants. *J Exp Med.* 2014;211(12):2361-72.
61. Bournazos S, DiLillo DJ, Ravetch JV. The role of Fc-FcγR interactions in IgG-mediated microbial neutralization. *J Exp Med.* 2015;212(9):1361-9.

62. DiLillo DJ, Ravetch JV. Differential Fc-Receptor Engagement Drives an Anti-tumor Vaccinal Effect. *Cell*. 2015;161(5):1035-45.
63. Hioe CE, Visciano ML, Kumar R, Liu J, Mack EA, Simon RE, et al. The use of immune complex vaccines to enhance antibody responses against neutralizing epitopes on HIV-1 envelope gp120. *Vaccine*. 2009;28(2):352-60.
64. Kumar R, Tuen M, Liu J, Nadas A, Pan R, Kong X, et al. Elicitation of broadly reactive antibodies against glycan-modulated neutralizing V3 epitopes of HIV-1 by immune complex vaccines. *Vaccine*. 2013;31(46):5413-21.
65. Villinger F, Mayne AE, Bostik P, Mori K, Jensen PE, Ahmed R, et al. Evidence for antibody-mediated enhancement of simian immunodeficiency virus (SIV) Gag antigen processing and cross presentation in SIV-infected rhesus macaques. *J Virol*. 2003;77(1):10-24.
66. Lambour J, Naranjo-Gomez M, Piechaczyk M, Pelegri M. Converting monoclonal antibody-based immunotherapies from passive to active: bringing immune complexes into play. *Emerg Microbes Infect*. 2016;5(8):e92.
67. Hua CK, Ackerman ME. Increasing the Clinical Potential and Applications of Anti-HIV Antibodies. *Front Immunol*. 2017;8:1655.
68. Zinkernagel RM, Bachmann MF, Kundig TM, Oehen S, Pirchet H, Hengartner H. On immunological memory. *Annu Rev Immunol*. 1996;14:333-67.
69. Pelegri M, Naranjo-Gomez M, Piechaczyk M. Antiviral Monoclonal Antibodies: Can They Be More Than Simple Neutralizing Agents? *Trends Microbiol*. 2015;23(10):653-65.
70. Pellegatta S, Eoli M, Frigerio S, Antozzi C, Bruzzzone MG, Cantini G, et al. The natural killer cell response and tumor debulking are associated with prolonged survival in recurrent glioblastoma patients receiving dendritic cells loaded with autologous tumor lysates. *Oncoimmunology*. 2013;2(3):e23401.
71. Gabrilovich D. Mechanisms and functional significance of tumour-induced dendritic-cell defects. *Nat Rev Immunol*. 2004;4(12):941-52.
72. Drakes ML, Stiff PJ. Regulation of Ovarian Cancer Prognosis by Immune Cells in the Tumor Microenvironment. *Cancers (Basel)*. 2018;10(9).
73. Halper-Stromberg A, Lu CL, Klein F, Horwitz JA, Bournazos S, Nogueira L, et al. Broadly neutralizing antibodies and viral inducers decrease rebound from HIV-1 latent reservoirs in humanized mice. *Cell*. 2014;158(5):989-99.

74. Chun TW, Murray D, Justement JS, Blazkova J, Hallahan CW, Fankuchen O, et al. Broadly neutralizing antibodies suppress HIV in the persistent viral reservoir. *Proc Natl Acad Sci U S A*. 2014;111(36):13151-6.
75. Horwitz JA, Halper-Stromberg A, Mouquet H, Gitlin AD, Tretiakova A, Eisenreich TR, et al. HIV-1 suppression and durable control by combining single broadly neutralizing antibodies and antiretroviral drugs in humanized mice. *Proc Natl Acad Sci U S A*. 2013;110(41):16538-43.
76. Shingai M, Nishimura Y, Klein F, Mouquet H, Donau OK, Plishka R, et al. Antibody-mediated immunotherapy of macaques chronically infected with SHIV suppresses viraemia. *Nature*. 2013;503(7475):277-80.
77. Pierson T, McArthur J, Siliciano RF. Reservoirs for HIV-1: mechanisms for viral persistence in the presence of antiviral immune responses and antiretroviral therapy. *Annu Rev Immunol*. 2000;18:665-708.
78. Estes JD, Kityo C, Ssali F, Swainson L, Makamdop KN, Del Prete GQ, et al. Defining total-body AIDS-virus burden with implications for curative strategies. *Nat Med*. 2017;23(11):1271-6.
79. Rothenberger MK, Keele BF, Wietgreffe SW, Fletcher CV, Beilman GJ, Chipman JG, et al. Large number of rebounding/founder HIV variants emerge from multifocal infection in lymphatic tissues after treatment interruption. *Proc Natl Acad Sci U S A*. 2015;112(10):E1126-34.
80. Danel C, Moh R, Minga A, Anzian A, Ba-Gomis O, Kanga C, et al. CD4-guided structured antiretroviral treatment interruption strategy in HIV-infected adults in west Africa (Trivacan ANRS 1269 trial): a randomised trial. *Lancet*. 2006;367(9527):1981-9.
81. Strategies for Management of Antiretroviral Therapy Study G, El-Sadr WM, Lundgren J, Neaton JD, Gordin F, Abrams D, et al. CD4+ count-guided interruption of antiretroviral treatment. *N Engl J Med*. 2006;355(22):2283-96.
82. Team DT. Fixed duration interruptions are inferior to continuous treatment in African adults starting therapy with CD4 cell counts < 200 cells/microl. *AIDS*. 2008;22(2):237-47.
83. Reynolds SJ, Kityo C, Hallahan CW, Kabuye G, Atwiine D, Mbamanya F, et al. A randomized, controlled, trial of short cycle intermittent compared to continuous antiretroviral therapy for the treatment of HIV infection in Uganda. *PLoS One*. 2010;5(4):e10307.
84. Firnhaber C, Azzoni L, Foulkes AS, Gross R, Yin X, Van Amsterdam D, et al. Randomized trial of time-limited interruptions of protease inhibitor-based antiretroviral therapy (ART) vs. continuous therapy for HIV-1 infection. *PLoS One*. 2011;6(6):e21450.
85. Lau JSY, Smith MZ, Lewin SR, McMahon JH. Clinical trials of antiretroviral treatment interruption in HIV-infected individuals. *AIDS*. 2019;33(5):773-91.

86. Routy JP, Boulassel MR, Nicolette CA, Jacobson JM. Assessing risk of a short-term antiretroviral therapy discontinuation as a read-out of viral control in immune-based therapy. *J Med Virol.* 2012;84(6):885-9.
87. Ananworanich J, Gayet-Ageron A, Le Braz M, Prasithsirikul W, Chetchotisakd P, Kiertiburanakul S, et al. CD4-guided scheduled treatment interruptions compared with continuous therapy for patients infected with HIV-1: results of the Staccato randomised trial. *Lancet.* 2006;368(9534):459-65.
88. Investigators ST, Fidler S, Porter K, Ewings F, Frater J, Ramjee G, et al. Short-course antiretroviral therapy in primary HIV infection. *N Engl J Med.* 2013;368(3):207-17.
89. Azzoni L, Foulkes AS, Papasavvas E, Mexas AM, Lynn KM, Mounzer K, et al. Improved treatment for primary HIV infection by interferon-alfa therapy? Does HCV treatment in HIV/HCV coinfectd patients help us to test this hypothesis? Reply to zur Wiesch and van Lunzen. *J Infect Dis.* 2013;208(2):363.
90. Jacobson JM, Routy JP, Welles S, DeBenedette M, Tcherepanova I, Angel JB, et al. Dendritic Cell Immunotherapy for HIV-1 Infection Using Autologous HIV-1 RNA: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *J Acquir Immune Defic Syndr.* 2016;72(1):31-8.
91. Gandhi RT, O'Neill D, Bosch RJ, Chan ES, Bucy RP, Shopis J, et al. A randomized therapeutic vaccine trial of canarypox-HIV-pulsed dendritic cells vs. canarypox-HIV alone in HIV-1-infected patients on antiretroviral therapy. *Vaccine.* 2009;27(43):6088-94.
92. Levy Y, Thiebaut R, Montes M, Lacabaratz C, Sloan L, King B, et al. Dendritic cell-based therapeutic vaccine elicits polyfunctional HIV-specific T-cell immunity associated with control of viral load. *Eur J Immunol.* 2014;44(9):2802-10.
93. Huiting ED, Gittens K, Justement JS, Shi V, Blazkova J, Benko E, et al. Impact of treatment interruption on HIV reservoirs and lymphocyte subsets in individuals who had initiated antiretroviral therapy during the early phase of infection. *J Infect Dis.* 2019.
94. Clarridge KE, Blazkova J, Einkauf K, Petrone M, Refsland EW, Justement JS, et al. Effect of analytical treatment interruption and reinitiation of antiretroviral therapy on HIV reservoirs and immunologic parameters in infected individuals. *PLoS Pathog.* 2018;14(1):e1006792.
95. Salantes DB, Zheng Y, Mampe F, Srivastava T, Beg S, Lai J, et al. HIV-1 latent reservoir size and diversity are stable following brief treatment interruption. *J Clin Invest.* 2018;128(7):3102-15.
96. Strongin Z, Sharaf R, VanBelzen DJ, Jacobson JM, Connick E, Volberding P, et al. Effect of Short-Term Antiretroviral Therapy Interruption on Levels of Integrated HIV DNA. *J Virol.* 2018;92(12).

97. Papasavvas E, Lada SM, Joseph J, Yin X, Liu Q, Azzoni L, et al. Analytical antiretroviral therapy interruption does not irreversibly change preinterruption levels of cellular HIV. *AIDS*. 2018;32(13):1763-72.
98. Stecher M, Claßen A, Klein F, Lehmann C, Gruell H, Platten M, et al. Systematic Review And Meta-Analysis of Treatment Interruptions in HIV-1 Infected Patients Receiving Antiretroviral Therapy: Implications for Future HIV Cure Trials. *Clinical Infectious Diseases*. 2019.
99. Hardy WD. Analytical Treatment Interruptions (ATIs) and HIV Cure Research Seizing the Opportunity, While Maintaining Safety and Respect. *Clinical Infectious Diseases*. 2019.
100. Michael NL. Clinical trial design: The nobility of randomization. *Sci Transl Med*. 2017;9(419).
101. Garcia F, Climent N, Guardo AC, Gil C, Leon A, Autran B, et al. A dendritic cell-based vaccine elicits T cell responses associated with control of HIV-1 replication. *Sci Transl Med*. 2013;5(166):166ra2.
102. Jacobson JM, Pat Bucy R, Spritzler J, Saag MS, Eron JJ, Jr., Coombs RW, et al. Evidence that intermittent structured treatment interruption, but not immunization with ALVAC-HIV vCP1452, promotes host control of HIV replication: the results of AIDS Clinical Trials Group 5068. *J Infect Dis*. 2006;194(5):623-32.
103. Rosenberg ES, Graham BS, Chan ES, Bosch RJ, Stocker V, Maenza J, et al. Safety and immunogenicity of therapeutic DNA vaccination in individuals treated with antiretroviral therapy during acute/early HIV-1 infection. *PLoS ONE*. 2010;5(5):e10555.
104. Schooley RT, Spritzler J, Wang H, Lederman MM, Havlir D, Kuritzkes DR, et al. AIDS clinical trials group 5197: a placebo-controlled trial of immunization of HIV-1-infected persons with a replication-deficient adenovirus type 5 vaccine expressing the HIV-1 core protein. *J Infect Dis*. 2010;202(5):705-16.
105. Henderson GE, Peay HL, Kroon E, Cadigan RJ, Meagher K, Jupimai T, et al. Ethics of treatment interruption trials in HIV cure research: addressing the conundrum of risk/benefit assessment. *J Med Ethics*. 2018;44(4):270-6.
106. Li JZ, Smith DM, Mellors JW. The need for treatment interruption studies and biomarker identification in the search for an HIV cure. *AIDS*. 2015;29(12):1429-32.
107. Henrich TJ, Hanhauser E, Marty FM, Sirignano MN, Keating S, Lee TH, et al. Antiretroviral-free HIV-1 remission and viral rebound after allogeneic stem cell transplantation: report of 2 cases. *Ann Intern Med*. 2014;161(5):319-27.

108. Hill AL, Rosenbloom DI, Goldstein E, Hanhauser E, Kuritzkes DR, Siliciano RF, et al. Real-Time Predictions of Reservoir Size and Rebound Time during Antiretroviral Therapy Interruption Trials for HIV. *PLoS Pathog.* 2016;12(4):e1005535.
109. Garner SA, Rennie S, Ananworanich J, Dube K, Margolis DM, Sugarman J, et al. Interrupting antiretroviral treatment in HIV cure research: scientific and ethical considerations. *J Virus Erad.* 2017;3(2):82-4.
110. Eyal N, Holtzman LG, Deeks SG. Ethical issues in HIV remission trials. *Curr Opin HIV AIDS.* 2018;13(5):422-7.
111. Edupuganti S, Mgodi N, Karuna S, Andrew P, Kochar N, deCamp A, et al. Antibody Mediated Prevention: Proof-of-Concept, Randomized, Blinded, Placebo-controlled Trials to Assess Safety and Efficacy of VRC01 to Prevent HIV-1 (P05.18LB). R4P, October 21-25, 2018, Madrid, Spain.
112. Andrew P, Andrasik M, Thommes E, deCamp A, Kochar N, Karg C, et al. Retention in the Ongoing AMP Trials of VRC01, a Broadly Neutralizing Antibody (bnAb) to Prevent HIV in Women, MSM & Transgender (TG) People (P05.17LB). R4P, October 21-25, 2018, Madrid, Spain.
113. Takuva S, Karuna S. HVTN 703/HPTN 081 & HVTN 704/HPTN 085 Antibody Mediated Prevention (AMP) Trials: Infusion Reactions Update. HVTN Sub-Saharan Africa Regional Meeting, 21-22 February 2018. Cape Town, South Africa.
114. Gilbert PB, Bosch RJ, Hudgens MG. Sensitivity analysis for the assessment of causal vaccine effects on viral load in HIV vaccine trials. *Biometrics.* 2003;59(3):531-41.
115. Benkeser D, Carone M, Gilbert PB. Improved estimation of the cumulative incidence of rare outcomes. *Stat Med.* 2018;37(2):280-93.
116. van der Laan MJ, Polley EC, Hubbard AE. Super learner. *Stat Appl Genet Mol Biol.* 2007;6:Article25.
117. Benkeser D, Hejazi N. *survtmle: Compute Targeted Minimum Loss-Based Estimates in Right-Censored Survival Settings [R package]*. 2018 [Available from: <https://cran.r-project.org/web/packages/survtmle/index.html>].
118. DeCamp A, Hraber P, Bailer RT, Seaman MS, Ochsenbauer C, Kappes J, et al. Global panel of HIV-1 Env reference strains for standardized assessments of vaccine-elicited neutralizing antibodies. *J Virol.* 2014;88(5):2489-507.
119. Seaman M, Janes H, Hawkins N, Grandpre L, Devoy C, Giri A, et al. Tiered Categorization of a Diverse Panel of HIV-1 Env Pseudoviruses for Assessment of Neutralizing Antibodies. *J Virol.* 2010;84(3):1439-52.

120. Tay MZ, Liu P, Williams LD, McRaven MD, Sawant S, Gurley TC, et al. Antibody-Mediated Internalization of Infectious HIV-1 Virions Differs among Antibody Isotypes and Subclasses. *PLoS Pathog.* 2016;12(8):e1005817.
121. Ackerman ME, Moldt B, Wyatt RT, Dugast AS, McAndrew E, Tsoukas S, et al. A robust, high-throughput assay to determine the phagocytic activity of clinical antibody samples. *J Immunol Methods.* 2011;366(1-2):8-19.
122. Tremblay CL, Hicks JL, Sutton L, Giguel F, Flynn T, Johnston M, et al. Antiretroviral resistance associated with supervised treatment interruptions in treated acute HIV infection. *AIDS.* 2003;17(7):1086-9.
123. Woodson C, Karim QA. A model designed to enhance informed consent: experiences from the HIV prevention trials network. *Am J Public Health.* 2005;95(3):412-9.
124. O'Connor AM, Wennberg JE, Legare F, Llewellyn-Thomas HA, Moulton BW, Sepucha KR, et al. Toward the 'tipping point': decision aids and informed patient choice. *Health Aff (Millwood).* 2007;26(3):716-25.
125. Pope TM. Certified Patient Decision Aids: Solving Persistent Problems with Informed Consent Law. *J Law Med Ethics.* 2017;45(1):12-40.
126. Steel JL, Gordon EJ, Dulovich M, Kingsley K, Tevar A, Ganesh S, et al. Transplant advocacy in the era of the human immunodeficiency virus organ policy equity act. *Clin Transplant.* 2018;32(7):e13309.
127. Durand CM, Segev D, Sugarman J. Realizing HOPE: The Ethics of Organ Transplantation From HIV-Positive Donors. *Ann Intern Med.* 2016;165(2):138-42.

Appendix A Sample informed consent form

Title: Antiretroviral analytical treatment interruption (ATI) to assess immunologic and virologic responses in participants who initiated ART in early HIV infection after having received VRC01 or placebo in HVTN 703/HPTN 081

Protocol number: HVTN 805/HPTN 093

Site: [Insert site name]

Thank you for your interest in our research study. Please read this consent form or ask someone to read it to you. If you decide to join the study, we will ask you to sign or make your mark on this form. We will offer you a copy to keep. We will ask you questions to see if we have explained everything clearly. You can also ask us questions about the study.

Research is not the same as treatment or medical care. The purpose of a research study is to answer scientific questions. These answers can help find new medicines, treatments, vaccines, and knowledge about how the human body works.

You are being invited to take part in this study because you got infected with HIV while you were enrolled in the AMP Study and started HIV treatment (also known as “ART” or anti-retroviral treatment) soon after the diagnosis, and because your viral load (the amount of HIV in your blood) has been kept at a very low level or “undetectable” for at least the past year. We will enroll people who got the study antibody and people who got placebo. Since the AMP Study is still blinded, we don’t know whether you got the study antibody or not.

We expect that between 16 and 61 people will join this study.

Key information

These are some of the things you should know about this study:

- One purpose of the study is to learn whether starting HIV treatment soon after diagnosis might help you control HIV after you interrupt your HIV treatment.
- A second purpose of the study is to learn whether having the AMP Study antibody in a person’s body might help the immune system control HIV better if that person gets HIV.
- A third purpose is to learn if it is safe for people to interrupt their HIV treatment in a carefully monitored research study.
- If you are eligible and choose to join, we will ask you to interrupt your HIV treatment.

- After you interrupt taking HIV treatment, you will have regular clinic visits to carefully track your HIV and your health. If your viral load goes up to 200 or higher, you will continue to stay off your HIV treatment, and we will check on your health more often.
- If your viral load goes up to 1,000 or more and stays that high for 4 weeks, or if your CD4 count drops below 350, or if you show any symptoms of HIV-related illness, you will restart taking HIV treatment. If your viral load at the end of 6 months is 200 or higher, you will restart taking HIV treatment. If your immune system maintains control of your HIV for the entire 6 months, you may choose to keep going without taking HIV treatment for as long as your viral load remains below 200 and you don't meet the other restart criteria within this study.
- Once you restart taking HIV treatment, you will have follow-up visits for up to a year to see how your body responds to restarting HIV treatment and to make sure that your viral load returns to very low or undetectable levels.
- At clinic visits we will give you physical exams, ask about your physical and mental health, and collect blood for laboratory tests.
- Interrupting your HIV treatment is an experimental procedure that is only used in carefully monitored research. It is not recommended as part of routine healthcare for people living with HIV. Healthcare guidelines recommend that people living with HIV take HIV treatment for the rest of their lives. Some of the risks when you interrupt HIV treatment include having a high viral load and acute retroviral syndrome, having a low CD4 count, transmitting HIV to your partner(s), developing HIV drug resistance, and having to switch to a new HIV treatment, which could have additional side effects. The study is designed to make it unlikely that any of these things will happen.
- There may also be risks we don't know about, even serious ones, including death.
- There is no direct benefit to you from being in the study.
- Whether to join this study is your choice. You do not have to join, and you are free to leave at any time. Your choice will not affect the care you receive at this clinic.

The rest of this form provides a more complete description of this study. Please read it carefully and ask questions at any time.

Why is this study being done?

Researchers know that when people interrupt their HIV treatment, their viral load usually goes up quickly, and they must restart taking their treatment. This quick increase does not always happen. Very rarely, for only a few people, it stays low, or even undetectable for many months.

We do not expect this study to benefit you directly, even if you received the study antibody in the AMP study. You should not expect to control the virus while you are off of your HIV treatment.

Researchers think that starting HIV treatment soon after being diagnosed with HIV may improve how the body's immune system responds to HIV and may keep the viral load low. Small studies in people have shown that people who start HIV treatment early are more likely to be able to control HIV after interrupting HIV treatment than people who started HIV treatment later. Researchers want to know how and why this happens.

Researchers also think that having an anti-HIV antibody like the one used in the AMP Study in the body at the time when HIV infection happens may improve how the body's immune system responds to HIV. This could happen even if the antibody didn't prevent HIV acquisition.

Animal studies suggest that having an anti-HIV antibody in the body at the time of HIV infection may help the immune system keep viral load low for a longer period of time without taking HIV treatment. Researchers want to find out if this will happen in people. And if it happens in people, researchers want to know how and why this happens.

1. We are doing this study to answer several questions.

- Are people who start HIV treatment soon after diagnosis likely to keep HIV under control after interrupting HIV treatment? If so, for how long?
- Are people who got the study antibody in the AMP Study more likely to keep HIV under control for a longer period of time without taking HIV treatment than people who got the placebo?
- Is it safe for people to interrupt taking their HIV treatment in this carefully monitored research study?
- Do people who got the study antibody in the AMP Study and got HIV develop and maintain an immune system that is more capable of controlling HIV viral load than people who got the placebo and get HIV?

- Do people who got the study antibody or placebo in the AMP Study and get HIV have a difference in the number of cells that are infected with HIV but do not reproduce before and after interrupting their HIV treatment?

2. To answer these questions, study participants will interrupt use of HIV treatment.

Interrupting HIV treatment is an experimental procedure that is only used in carefully monitored research. It is not recommended as part of routine healthcare for people living with HIV. Healthcare guidelines recommend that people living with HIV take HIV treatment for the rest of their lives. We will tell you more about the risks of interrupting your HIV treatment in Section 19 of this form. During the time that you do not take your HIV treatment we will check your health and HIV frequently to minimize the chance of harm to you. If your viral load goes up, your CD4 count goes down, you show signs of HIV-related illness, or you and/or your primary healthcare worker decide to, you will restart taking your HIV treatment right away.

Joining the study

3. It is completely up to you whether or not to join the study.

Take your time in deciding. If it helps, talk to people you trust, such as your primary HIV care provider, friends or family. We will be happy to help you explain this study to anyone you wish. If you decide not to join this study, or if you leave it after you have joined, your other care at this clinic and the benefits or rights you would normally have will not be affected.

You cannot be in this study while you are in another study where you get a study product. Being in more than one study may not be safe.

If you choose not to join this study, you may be able to join another study.

4. If you want to join the study, we will screen you to see if you are eligible.

Screening involves a physical exam, checking your mental health, an HIV test, and asking about your health history. A physical exam may include, but is not limited to:

- Checking your weight, temperature and blood pressure,
- Looking in your mouth and throat,
- Listening to your heart and lungs, and
- Feeling your abdomen (stomach and liver).

We will also do blood and urine tests. These tests tell us about some aspects of your health, such as how healthy your kidneys, liver, and immune system are. We will also test you for hepatitis B, hepatitis C, tuberculosis (TB), gonorrhea, chlamydia, syphilis, and Trichomonas. We will ask you about medications you are taking. If you can become pregnant, we will test you for pregnancy. If you are pregnant, you cannot join the study.

We will communicate with your primary healthcare worker to coordinate your care during this study. We will check to make sure your HIV has been well controlled by your HIV treatment for at least one year.

We will ask if you are willing to use condoms every time you have sex for an extended period of time during the study, until you restart your HIV treatment and your viral load returns to being very low or undetectable. This is to protect your partner(s).

We will review the screening results with you. The screening results may show you are not eligible to join the study, even if you want to.

5. If we find that you have a health problem during screening or during the study, we will tell you about the care that we can give here for free.

For the care that we cannot give, we will explain how we will help you get care elsewhere. We will not pay for care for health problems that are unrelated to the study.

Site: Appendix B, Approved contraception methods (for sample informed consent form), contains supplemental information. If you want to include Appendix B in this consent form, paste it below.

6. If you can become pregnant, you must agree to use effective contraception to join this study.

Current HIV treatment guidelines recommend using HIV treatment during pregnancy to prevent transmission to the baby. Since you will interrupt your HIV treatment in this study, you should not become pregnant. In addition to using condoms every time you have sex to prevent HIV transmission to your partner, we will ask you to use contraception as well. Effective contraception can include an IUD or hormone-based contraception.

If you become pregnant or start breastfeeding before you interrupt your HIV treatment, you will have to leave the study immediately.

If you become pregnant or start breastfeeding after you interrupt your HIV treatment, you will have to restart taking it, but you can remain in the study. If you become pregnant or start breastfeeding after you have restarted your HIV treatment, you will continue taking it. In both these situations, we would like you

to come for clinic visits for up to 6 months after your HIV viral load returns to very low or undetectable.

The study staff would like to obtain information from you about the outcome of the pregnancy (even if it is after your participation in the study ends). If you are taking your HIV treatment when you become pregnant, your pregnancy will be reported to an international database that collects information about pregnancies in people using HIV treatments. This report will not use your name or other information that could be used to identify you.

Being in the study

If you meet the study requirements and want to join, here is what will happen.

7. We will make sure you are on the right HIV treatment for this study.

You may have to change some of the HIV treatment you are taking. If you are taking HIV treatment that stays in the blood for a long time even after you stop taking it, we will have you switch to a different HIV treatment for at least one month before you interrupt all of your HIV treatment. We will provide this new treatment to you for free. This is to reduce the already small chance that you may develop drug resistance. Drug resistance means the HIV treatment you have been taking might not control your HIV when you restart it. We will check on you after about 2 weeks to see how you are doing on the new HIV treatment. If you have any problems after starting the new HIV treatment, please let us know.

We will ask you to come to the clinic about 4 weeks after starting the new HIV treatment to see if it is controlling your HIV. If it is not, we will ask you to come to the clinic weekly for up to 12 weeks (about 3 months) until your HIV is controlled.

If the new HIV treatment does not bring your HIV viral load to very low or undetectable within about 3 months, you will not be able to continue with the rest of the study.

8. If you join the study, we will collect some basic information.

We will record your medical history and give you a physical examination, including checking your weight and vital signs. We will ask about other medications you are taking and about any illnesses you may have. We will also collect blood and urine samples, and if you can become pregnant, we will give you a pregnancy test. We will test you for gonorrhea and chlamydia using urine or a cervical/vaginal swab. We will test you for syphilis using a blood sample and for Trichomonas using a cervical/vaginal swab.

9. In Part 1 of the study, you will interrupt your HIV treatment.

After you interrupt your HIV treatment, for the first 8 weeks (about 2 months) you will visit the clinic every week (8 visits). For the next 16 weeks (months 2-6), you will visit the clinic every 2 weeks (8 visits). For the next 6 months, if you are still in Part 1 of the study, you will visit the clinic about every month (7 visits). One reason for these frequent clinic visits is to carefully track your HIV and your health.

At these visits we will ask about anything new in your medical history and give you a brief physical exam. We will collect blood. When we collect blood, the amount will depend on the lab tests we need to do. It will be some amount between 6 mL and 130 mL (a little more than 1 teaspoon and a little less than ½ cup). Your body will make new blood to replace the blood we take out.

We will check to make sure you are using condoms for all sexual activity.

If you can become pregnant, we will check to make sure you are using effective contraception. We will also give you a pregnancy test.

We will test you for gonorrhea and chlamydia at some visits using urine or a cervical/vaginal swab. At these times we will also test you for Trichomonas using a cervical/vaginal swab.

Some of the blood we collect will be used for lab tests to check your general health (including syphilis testing) and to check your immune system. Estrogen and progesterone may affect the immune system, so we will check the amount of those hormones in your blood. We will also check to see if you still have any HIV treatment in your blood.

We will review the results of these procedures tests with you at your next visit, or sooner if necessary. If any of the results are important to your health, we will tell you.

Some of the blood collected at each visit will be used to test your viral load and your CD4 count. If your viral load goes up to 200 or higher, you will move to Part 2 of the study.

If any of the following happens, you will restart your HIV treatment as soon as possible and move to Part 3 of the study.

- Your CD4 count drops below 350,
- You show any symptoms of HIV-related illness,
- You decide to restart your HIV treatment, or

- Your primary healthcare worker or the doctor at the study clinic decides that you should restart your HIV treatment.

If none of these things happens, a person can stay in Part 1 for a year or longer. Most people will be in Part 1 of the study for less than 6 months.

If you meet the restart criteria but choose not to restart ART, we will ask you to continue Part 1 clinic visits. We may need to see you for additional visits to check your safety. If you restart your HIV treatment you will move to Part 3.

10. Here is what will happen in Part 2 of the study.

In Part 2, you will continue to stay off your HIV treatment. When you start Part 2, for the first 8 weeks (about 2 months), you will visit the clinic every week (9 visits). For the next 28 weeks (months 2-8), you will visit the clinic every 2 weeks (14 visits). After that, if you still have not met the criteria to restart your HIV treatment, for the next 4 months, you will visit the clinic once a month (4 visits). One reason for these frequent clinic visits is to carefully track your HIV and your health.

At these visits we will ask about anything new in your medical history. At the first visit in Part 2, we will give you a complete physical exam. At all other visits in Part 2 we will give you a brief physical exam. We will collect blood. When we collect blood, the amount will depend on the lab tests we need to do. It will be some amount between 6 mL and 120 mL (a little more than 1 teaspoon and a little less than ½ cup). Your body will make new blood to replace the blood we take out.

We will check to make sure you are using condoms for all sexual activity.

If you can become pregnant, we will check to make sure you are using effective contraception. We will also give you a pregnancy test.

We will test you for gonorrhea and chlamydia at some visits using urine or a cervical/vaginal swab. At these times we will also test you for Trichomonas using a cervical/vaginal swab.

Some of the blood we collect will be used for lab tests of your general health (including syphilis testing) and to check your immune system. Estrogen and progesterone may affect the immune system, so we will check the amount of those hormones in your blood. We will also check to see if you have any HIV treatment in your blood.

We will review the results of these procedures and tests with you at your next visit, or sooner if necessary. If any of the results are important to your health, we will tell you.

Some of the blood collected at each visit will be used to test your viral load and your CD4 count.

During the first 24 weeks (about 6 months) in Part 2, if any of the following happens, you will restart your HIV treatment as soon as possible and move to Part 3 of the study.

- Your viral load goes up to 1,000 or higher and stays there for 4 weeks without dropping very quickly,
- Your CD4 count drops below 350,
- You show any symptoms of HIV-related illness,
- You decide to restart your HIV treatment, or
- Your primary HIV care provider or the doctor at the study clinic decides that you should restart your HIV treatment.

After 24 weeks (6 months), if any of the following happens, you will restart your HIV treatment as soon as possible and move to Part 3 of the study.

- Your viral load is 200 or higher,
- Your CD4 count drops below 350,
- You show any symptoms of HIV-related illness,
- You decide to restart your HIV treatment, or
- Your primary healthcare worker or the doctor at the study clinic decides that you should restart your HIV treatment.

If you do not meet these restart criteria, you can stay in Part 2 of the study for as long as your viral load stays below 200. Most people will be in Part 2 for a few weeks to a few months. Some people may stay in Part 2 for a year or longer. If you meet the restart criteria but choose not to restart ART, we will ask you to continue Part 2 clinic visits. We may need to see you for additional visits to check your safety. If you restart your HIV treatment you will move to Part 3.

11. After you restart taking HIV treatment, you will have follow-up clinic visits for about another year. This is Part 3 of the study.

This follow-up period is to see how your body responds to restarting your HIV treatment and to make sure that your HIV returns to very low or undetectable levels. For most people, we expect that your viral load will drop within about 12

weeks (about 3 months). If it does not, we will ask you to come to the clinic every 2 weeks until your HIV is controlled.

Once you restart taking HIV treatment, for the first 12 weeks (about 3 months) you will have clinic visits every 2 weeks (7 visits). For the next 16 weeks (about 4 months), you will have clinic visits once a month (4 visits). For the next 24 weeks (about 6 months) you will have 2 visits scheduled 3 months apart. At these visits we will ask about anything new in your medical history and give you a brief physical exam. We will collect blood. When we take blood, the amount will depend on the lab tests we need to do. It will be some amount between 6 mL and 225 mL (a little more than 1 teaspoon and a little less than 1 cup).

At some visits, we will test you for gonorrhea and chlamydia using urine or a cervical/vaginal swab. At these times we will also test you for *Trichomonas* using a cervical/vaginal swab. We will also do a blood test for syphilis.

We will review the results of these procedures and tests with you at your next visit, or sooner if necessary. If any of the results are important to your health, we will tell you.

You will still need to use condoms every time you have sex until we have confirmed that your viral load has returned to very low or undetectable levels. If you can become pregnant, we will check to make sure you are using effective contraception until we have confirmed that your viral load has returned to very low or undetectable levels. We will also give you a pregnancy test at some visits.

12. If you stay in Part 1 or Part 2 of the study for more than a year, you will have visits every 3 months.

At these visits we will ask about anything new in your medical history and give you a brief physical exam. We will collect blood. Some of that blood gets used for viral load and CD4 testing.

At some visits, we will test you for gonorrhea and chlamydia using urine or a cervical/vaginal swab. At these times we will also test you for *Trichomonas* using a cervical/vaginal swab. At these same visits, we will test you for syphilis using blood samples.

We will review the results of these tests with you at your next visit, or sooner if necessary. If any of the results are important to your health, we will tell you.

We will continue to check to make sure you are using condoms every time you have sex.

Also, if you can become pregnant, we will check to make sure you are using effective contraception. We will also give you a pregnancy test at each visit.

If you meet the restart criteria at any time, you will restart your HIV treatment and move to Part 3 of the study as described above in Section 11. If you choose not to restart your treatment, we will strongly urge you to continue clinic visits on the same schedule you have been on. We may need to see you for additional visits to check your safety. If you restart your HIV treatment you will move to Part 3.

13. Most participants will be in the study about 13-18 months.

This depends on whether you need to switch your HIV treatment when you enroll and how long it takes before you meet the criteria to restart your HIV treatment. If your immune system controls your HIV for a long time without treatment, then you might be in the study longer. The longest we expect anyone to be in the study is about 3 years.

14. We will test your samples to see how your immune system is functioning.

We will send your samples (without your name) to labs approved by the HVTN and the HPTN for this study, which are located in the United States and South Africa. In rare cases, some of your samples may be sent to labs approved by the HVTN and HPTN in other countries for research related to this study.

Some of the blood we collected previously in the AMP Study or that we collect in this study might be used for genetic testing in this study. Your genes are passed to you from your birth parents. They affect how you look and how your body works. Differences in genes can explain why some people get a disease and others don't. The genetic testing that might be done in this study involves only some of your genes related to how your immune system works to fight HIV. The testing will not involve all of your genes (your genome). We want to understand whether your genes affect your ability to control HIV in combination with whether or not you got the AMP Study antibody.

In some cases, researchers may take cells from your samples and grow more of them over time, so that your samples can continue to contribute to this study.

These tests done on your samples are for research purposes, not to check your health. The labs will not give the results to you or this clinic because their tests are not approved for use in making health care decisions. These labs are only approved to do research tests.

When your samples are no longer needed for this study, the HVTN will continue to store them.

Site: Delete next section if using separate consent for use of samples and information in other studies.

15. When samples are no longer needed for this study, the HVTN and HPTN may want to use them in other studies and share them with other researchers.

These samples are called “extra samples”. The HVTN and HPTN will only allow your extra samples to be used in other studies if you agree to this. You will mark your decision at the end of this form. If you have any questions, please ask.

Do I have to agree? No. You are free to say yes or no, or to change your mind after you sign this form. At your request, the HVTN and HPTN will destroy all extra samples that they have. Your decision will not affect your being in this study or have any negative consequences here.

Where are the samples stored? Extra samples are stored in a secure central place called a repository. Your samples will be stored in the HVTN repository in the South Africa.

How long will the samples be stored? There is no limit on how long your extra samples will be stored. *[Site: Revise the previous sentence to insert limits if your regulatory authority imposes them.]*

Will I be paid for the use of my samples? No. Also, a researcher may make a new scientific discovery or product based on the use of your samples. If this happens, there is no plan to share any money with you. The researcher is not likely to ever know who you are.

Will I benefit from allowing my samples to be used in other studies? Probably not. Results from these other studies are not given to you, this clinic, or your doctor. They are not part of your medical record. The studies are only being done for research purposes.

Will the HVTN or HPTN sell my samples and information? No, but the HVTN and HPTN may share your samples with other researchers. Once we share your samples and information, the HVTN and HPTN may not be able to get them back.

How do other researchers get my samples and information? When a researcher wants to use your samples and information, their research plan must be approved by the HVTN and HPTN. Also, the researcher’s institutional review board (IRB) or ethics committee (EC) will review their plan. *[Site: If review by your institution’s IRB/EC/RE is also required, insert a sentence stating this.]* IRBs/ECs protect the rights and well-being of people in research. If the research plan is approved, the HVTN will send your samples to the researcher’s location.

What information is shared with HVTN, HPTN, or other researchers? The samples and information will be labeled with a code number. The key to the code will stay at this clinic. It will not be shared with the HVTN, other researchers or with anyone else who does not need to know your name. Your name will not be

part of the information. However, some information that The HVTN and HPTN share may be personal, such as your race, ethnicity, sex, health information from the study, and HIV status. The HVTN and HPTN may share information about the study product you received and how your body responded to the AMP study product.

What kind of studies might be done with my extra samples and information? The studies will be related to HIV, vaccines, monoclonal antibodies, the immune system, and other diseases.

Researchers may also do genetic testing on your samples.

In some cases, researchers may take cells from your samples and grow more of them over time, so that they can continue to do research with them.

If you agree, your samples could also be used for genome wide studies. In these studies, researchers will look at all your genes (your genome). The researchers compare the genomes of many people, looking for common patterns of genes that could help them understand diseases. The researchers may put the information from the genome-wide studies into a protected database so that other researchers can access it. Usually, no one would be able to look at your genome and link it to you as a person. However, if another database exists that also has information on your genome and your name, someone might be able to compare the databases and identify you. If others found out, it could lead to discrimination or other problems. The risk of this happening is extremely small.

Who will have access to my information in studies using my extra samples?

People who may see your information are:

- Researchers who use your extra samples and information for other research,
- Government agencies that fund or monitor the research using your extra samples and information,
- Any regulatory agency that reviews clinical trials,
- The researcher's Institutional Review Board or Ethics Committee, or
- The people who work with the researcher/

All these people will do their best to protect your information. The results of any new studies that use your extra samples and information may be published. No publication will use your name or identify you personally.

16. We will reimburse you [Site: Insert compensation] for each study visit you complete.

This amount is to cover the costs of [Site: Insert text]

Site: Insert any costs to participants (eg, contraception costs for participants who could become pregnant).

You do not have to pay anything to be in this study.

17. We will do our best to protect your private information.

Your study records and samples will be kept in a secure location. We will label all of your samples and most of your records with a code number, not your name or other personal information. However, it is possible to identify you, if necessary. We will not share your name with the lab that does the tests on your samples, or with anyone else who does not need to know your name.

Clinic staff will have access to your study records. Your records may also be reviewed by groups who watch over this study to see that we are protecting your rights, keeping you safe, and following the study plan. These groups include:

- The US National Institutes of Health and its study monitors,
- [Insert name of local IRB/EC] ,
- [Insert name of local and/or national regulatory authority as appropriate],
- Any regulatory agency that reviews clinical trials,
- The Dale and Betty Bumpers Vaccine Research Center and people who work for them,
- The HVTN, HPTN and people who work for them,
- The US National Institute of Allergy and Infectious Diseases Data and Safety Monitoring Board, and
- The US Office for Human Research Protections.

All reviewers will take steps to keep your records private.

We cannot guarantee absolute privacy. If you are found to have a medical condition that we are required to report by law, then some of your information may be shared. At this clinic, we have to report the following information:

Site: Include any public health or legal reporting requirements. Bulleted examples should include all appropriate cases (reportable communicable disease, risk of harm to self or others, etc.). If your site does not have public health or legal reporting requirements, you may delete the last sentence in the paragraph above, along with the bullets below.

- [Item 1]
- [Item 2]
- [Item 3]

The results of this study may be published. No publication will use your name or identify you personally.

We may share information from the study with other researchers. We will not share your name or information that can identify you.

A description of this clinical trial will be available on <http://www.ClinicalTrials.gov>. This website will not include information that can identify you. At most, the website will include a summary of the results. You can search this website at any time.

18. We may take you out of the study even if you want to stay in it.

This may happen if:

- You move away from the study site and transferring to another study site is not possible,
- Clinic staff cannot contact you,
- We think, or your primary HIV care provider thinks, that staying in the study might harm you,
- You do not follow study instructions,
- You enroll in a different research study where you get a study product,
- You become pregnant or are breastfeeding,
- You tell us you intend to get pregnant or begin breastfeeding while you are not taking your HIV treatment, or
- The study is stopped for any reason.

Risks

19. There are risks to being in this study.

This study is designed to minimize the risks of interrupting your HIV treatment. This depends on you following the instructions from the clinic staff and attending all your study visits.

This section describes the risks we know about. There may also be risks we don't know about, even serious ones. We will tell you if we learn anything new that may affect your willingness to stay in the study.

Risks of interrupting your HIV treatment

Staying off HIV treatment is not recommended for routine healthcare for people living with HIV. Interrupting HIV treatment is an experimental procedure. It can increase the risk of serious health complications, including death. This section describes the risks we know about. There may also be risks we don't know about, even serious ones.

High viral load

At some point after you interrupt your HIV treatment, your viral load will probably rise. During this time, you probably will not have any symptoms. However, it is possible that you could have symptoms of what is called “acute retroviral syndrome.” We think that having this syndrome is rare because you started taking HIV treatment soon after your HIV diagnosis. The symptoms could include:

- Fever,
- Weight loss,
- Sore throat,
- Swollen lymph nodes,
- Headache,
- Body or joint aches,
- Tiredness,
- Night sweats,
- Oral yeast infection,

- Diarrhea, or
- Rash.

If you have any of these symptoms, tell the study staff right away. These symptoms should go away or become less severe once you restart your HIV treatment.

If your viral load increases, this could lead to inflammation. High levels of inflammation over many years are associated with heart disease and other medical conditions. Starting your HIV treatment again should reduce inflammation.

Low CD4 count

If your viral load rises, your CD4 count could drop. This could increase your risk for other HIV-related illnesses. In about 60 people studied to date, there has not been a significant difference in CD4 counts before and 6-12 months after interrupting HIV treatment. However, it is possible that your CD4 count might not return to its original level, even after you restart taking your HIV treatment.

Transmitting HIV to your partner(s)

While your viral load is controlled, the risk that you could transmit HIV to a sexual partner is very low, known as U = U (undetectable = untransmittable). Once you interrupt HIV treatment that risk gets higher.

That is why you must use condoms for all sexual activity during the time that you are not taking your HIV treatment.

If your sexual partners do not have HIV, they should get regular HIV testing until you have restarted your HIV treatment and we have confirmed that your viral load has returned to very low or undetectable levels. Testing for your partners is not part of this study.

Site: Adjust the next sentence as appropriate for your site.

If your partners are interested, we can tell you about a program we have for providing pre-exposure prophylaxis (PrEP) to them during the time that you are in the study. You are also welcome to bring partners to the clinic for education and risk reduction counseling.

Transmitting HIV to your baby

If you become pregnant, the risk of transmitting HIV to your baby gets higher if you are not taking your HIV treatment. That is why you must keep using contraception during the time when you are not taking HIV treatment until you restart treatment and your viral load returns to very low or undetectable. If you

become pregnant, it is extremely important to take your HIV treatment throughout the entire pregnancy.

Developing HIV drug resistance

As we described in Section 7 above, when you interrupt HIV treatment, the HIV in your body might mutate in ways that make it resistant to the HIV treatment you have been taking. This means the treatment you were taking may not control your HIV when you start taking it again. To make this less likely, we may ask you to switch to a different HIV treatment before beginning the study.

After you restart your HIV treatment, you will have very frequent viral load tests. If your viral load is not returning to its original level within 8-12 weeks of restarting your HIV treatment, we will test to see if the HIV is resistant to your HIV treatment regimen. If resistance is found, we will recommend more effective HIV treatment to you and your primary healthcare worker.

Side effects from new HIV treatment or non-study medications:

If you need to switch your HIV treatment, there may be some new side effects from the new HIV treatment. We or your primary healthcare worker can explain those side effects to you.

There may also be a risk of serious or even life-threatening side effects from non-study medications taken during the study. For your safety, tell the study doctor or nurse at the study clinic about all medications you are taking before you start the study, and also before starting using any new medications during the study.

Routine medical procedures:

In this study, we will take blood. This can cause bruising, pain, fainting, soreness, redness, swelling, itching, a sore, bleeding, and (rarely) muscle damage or infection where the needle was inserted. In some people, taking blood can cause a low blood cell count (anemia), making you feel tired.

Risks of disclosure of your personal information:

We will take several steps to protect your personal information. Although the risk is very low, it is possible that your personal information could be given to someone who should not have it. If that happened, you could face discrimination, stress, and embarrassment. We can tell you more about how we will protect your personal information if you would like.

Risks of other people learning your HIV status:

We will keep your participation in this study and your HIV status confidential. However, if others learn that you are in a study for people living with HIV, you

could face discrimination, stress, embarrassment, stigma, or intimate partner violence.

Risk of limitations to research participation:

As a result of being in this study, you may not be eligible to participate in other HIV treatment or cure studies that require you to have an undetectable viral load for a period of time. After you restart taking your HIV treatment and your viral load becomes undetectable again, this limitation will go away over time.

Risks of genetic testing:

It is unlikely, but the genetic tests that might be done on your samples could show you may be at risk for certain diseases. If others found out, it could lead to discrimination or other problems. However, it is almost impossible for you or others to know your test results from the genetic testing. The results are not part of your study records and are not given to you.

Benefits

20. We do not expect the study to benefit you directly.

However, information learned from this study may help others who have HIV and being in the study might still help you in some ways. The lab tests and physical exams that you get while in this study might detect health problems you don't yet know about. You may also learn whether you are one of the rare people who can control HIV without using HIV treatment. This study may help in the search for a vaccine or other ways to prevent HIV.

Your rights and responsibilities

21. If you join the study, you have rights and responsibilities.

You have many rights that we will respect. You also have responsibilities. We list these in the Participant's Bill of Rights and Responsibilities. We will give you a copy of it.

Leaving the study

22. Tell us if you decide to leave the study.

You are free to leave the study at any time and for any reason. Your care at this clinic and your legal rights will not be affected, but it is important for you to let us know.

We will ask you to come back to the clinic one last time for a physical exam, and we may ask to take some blood and urine samples. We will also ask about any

personal problems or benefits you have experienced from being in the study. We will counsel you about the importance of taking your HIV treatment and following up with your primary healthcare worker. We believe these steps are important to protecting your health, but it is up to you whether to complete them.

Injuries

Sites: Approval from HVTN Regulatory Affairs (at vtn.core.reg@hvtvn.org) is needed for any change (other than those that the instructions specifically request or those previously approved by HVTN Regulatory Affairs) to the boxed text

23. If you get sick or injured during the study, contact us immediately.

Your health is important to us. *(Sites: adjust the following 2 sentences if applicable to the care available at your site)* We will tell you about the care that we can give here. For the care that we cannot provide, we will explain how we will help you get care elsewhere.

If you become sick or injured in this study, the HVTN has a process to decide if it is related to the procedures. If it is, we call it a study-related injury. There are funds to pay for treatment of study-related injuries if certain conditions are met.

Sites: adjust the language in this paragraph so it is applicable to your site. Note that the ABPI guidelines apply to South Africa only. In this study, our clinic has insurance to cover your medical treatment in the case of a study-related injury. We will follow the Association of the British Pharmaceutical Industry guidelines for payment of study-related injury. We can give you a copy of these guidelines. In rare cases, the insurance funds may not be enough.

Some injuries are not physical. For example, you might be harmed emotionally by being in a study where you interrupt your HIV treatment. Or you might lose wages because you cannot go to work. However, there are no funds to pay for these kinds of injuries, even if they are study related.

You may disagree with the decision about whether your injury is study related. If you wish, the HVTN will ask independent experts to review the decision. You always have the right to use the court system if you are not satisfied.

Questions

24. If you have questions or problems at any time during your participation in this study, use the following important contacts.

If you have questions about this study, contact
[name or title and telephone number of the investigator or other study staff].

If you have any symptoms that you think may be related to this study, contact [name or title and telephone number of the investigator or other study staff].

This study has been reviewed and approved by a committee called the [name of local IRB/EC]. If you have questions about your rights as a research participant, or problems or concerns about how you are being treated in this study, contact [name or title and telephone number of person on IRB/EC] , at the committee.

Paragraph for South African sites. The study has been structured in accordance with the Declaration of Helsinki (last updated October 2013) which deals with the recommendations guiding doctors in biomedical research involving human participants, the Ethics in Health Research: Principles, Structures and Processes Second Edition 2015, and Guidelines for Good Practice in the Conduct of Clinical Trials in Human Participants in South Africa. We can provide you with copies of these guidelines if you wish to review them.

If you want to leave this study, contact [name or title and telephone number of the investigator or other study staff].

Remainder of section for South African sites only.

You can reach a study staff member 24 hours a day at [telephone number].

If you have questions about this trial you should first discuss them with your doctor or the ethics committee (contact details as provided on this form). After you have consulted your doctor or the ethics committee and if they have not provided you with answers to your satisfaction, you should write to the South African Healthcare Products Regulatory Authority (SAHPRA) at:

The Acting Chief Executive Officer
South African Health Products Regulatory Authority
Private Bag X828
PRETORIA
0001

Tel: (012) 842 7582/3

e-mail: portia.nkambule@sahpra.org.za

Your permissions and signature

25. In Section 15 of this form, we told you about possible other uses of your extra samples and information, outside this study. Please choose only one of the options below and write your initials or make your mark in the box next to it. Whatever you choose, the HVTN keeps track of your decision about how

your samples and information can be used. You can change your mind after signing this form.

☐

I allow my extra samples and information to be used for other studies related to HIV, vaccines, monoclonal antibodies, the immune system, and other diseases. This may include genetic testing and keeping my cells growing over time.

OR

☐

I agree to the option above *and* also to allow my extra samples and information to be used in genome wide studies.

OR

☐

I do not allow my extra samples to be used in any other studies. This includes not allowing genetic testing, growing more of my cells, or genome wide studies.

26. If you agree to join this study, you will need to sign or make your mark below. Before you sign or make your mark on this consent form, make sure of the following:

- You have read this consent form, or someone has read it to you.
- You feel that you understand what the study is about and what will happen to you if you join. You understand what the possible risks and benefits are.
- You have had your questions answered and know that you can ask more.
- You agree to join this study.

You will not be giving up any of your rights by signing this consent form.

Participant's name (print)	Participant's signature or mark	Date	Time
Clinic staff conducting consent discussion (print)	Clinic staff signature	Date	Time

For participants who are unable to read or write, a witness should complete the signature block below:

Witness's name (print)

Witness's signature

Date

Time

*Witness is impartial and was present for the entire discussion of this consent form.

Appendix B Approved birth control methods (for sample informed consent form)

If you can become pregnant, you must agree to use effective contraception from 21 days before interrupting your HIV treatment until you start taking HIV treatment again and your viral load drops to undetectable.

You must use condoms every time you have sex to prevent HIV transmission to your partner. You must also use one of the following contraceptive methods:

- Birth control drugs that prevent pregnancy—such as pills, patches, vaginal rings, injectables, or an implant under the skin; or
- Intrauterine device (IUD).

You do not have to use birth control if:

- You have been diagnosed with early menopause, with no menstrual periods for one year;
- You have had a hysterectomy (your uterus removed);
- You have had your ovaries removed;
- You have a tubal ligation (your “tubes tied”) or confirmed successful placement of a product that blocks the fallopian tubes.

Appendix C Sample consent form for use of samples and information in other studies

Title: Antiretroviral analytical treatment interruption (ATI) to assess immunologic and virologic responses in participants who initiated ART in early HIV infection after having received VRC01 or placebo in HVTN 703/HPTN 081

Protocol number: HVTN 805/HPTN 093

Site: [Insert site name]

When samples are no longer needed for this study, the study sponsors want to keep them for use in other studies by HVTN, HPTN, or other researchers. We will call these “extra samples.”

This form gives you information so you can decide if you want your extra samples and information used in other studies. You will mark your decision at the end of the form. If you have any questions, please ask.

Key information

These are some of the things you should know about the use of your samples and information for other studies:

- The extra samples will be labeled with a code number and some personal information. They will not be labeled with your name. The extra samples are stored in a secure place. At your request, the HVTN and HPTN will destroy all your extra samples. You can still join the main study even if you do not agree to the use of your extra samples in other studies.
- Researchers may do genetic testing on your samples, which could include genome wide studies. It is unlikely, but these tests could show you may be at risk for certain diseases. In the very unlikely event that others found out, this could lead to discrimination or other problems.
- You will not be paid or otherwise benefit from allowing your extra samples to be used in other studies.

The rest of this form gives more information about use of your extra samples for other studies. Please read it carefully.

1. Do I have to agree?

No. You are free to say yes or no, or to change your mind after you sign this form. At your request, the HVTN and HPTN will destroy all extra samples that they have. Your decision will not affect your being in this study or have any negative consequences here.

2. Where are the samples stored?

Extra samples are stored in a secure central place called a repository. Your samples will be stored in the HVTN repository in South Africa.

3. How long will the samples be stored?

There is no limit on how long your extra samples will be stored. *[Site: Revise the previous sentence to insert limits if your regulatory authority imposes them.]*

4. Will I benefit from allowing my samples to be used in other studies?

We do not expect this to benefit you directly. Results from these other studies are not given to you, this clinic, or your doctor. They are not part of your medical record. The studies are only being done for research purposes.

5. Will I be paid for the use of my samples?

No. Also, a researcher may make a new scientific discovery or product based on the use of your samples. If this happens, there is no plan to share any money with you. The researcher is not likely to ever know who you are.

6. Will the HVTN or HPTN sell my samples and information?

No, but the HVTN and HPTN may share your samples with other researchers. Once we share your samples and information, we may not be able to get them back.

7. How do other researchers get my samples and information?

When a researcher wants to use your samples and information, their research plan must be approved by the HVTN and HPTN. Also, the researcher's institutional review board (IRB) or ethics committee (EC) will review their plan. *[Site: If review by your institution's IRB/EC/RE is also required, insert a sentence stating this.]* IRBs/ECs protect the rights and well-being of people in research. If the research plan is approved, the HVTN and HPTN will send your samples to the researcher's location.

8. What information is shared with HVTN, HPTN, or other researchers?

The samples and information will be labeled with a code number. The key to the code will stay at this clinic. It will not be shared with the HVTN, other researchers or with anyone else who does not need to know your name. Your name will not be part of the information. However, some information that the HVTN and HPTN share may be personal, such as your race, ethnicity, gender, health information from the study, and HIV status. The HVTN and HPTN may

share information about the study product you received and how your body responded to the AMP study product.

9. What kind of studies might be done with my extra samples and information?

The studies will be related to HIV, vaccines, monoclonal antibodies, the immune system, and other diseases.

Researchers may also do genetic testing on your samples.

In some cases, researchers may take cells from your samples and grow more of them over time, so that they can continue to contribute to this study.

If you agree, your samples could also be used for genome wide studies. In these studies, researchers will look at all of your genes (your genome). The researchers compare the genomes of many people, looking for common patterns of genes that could help them understand diseases. The researchers may put the information from the genome-wide studies into a protected database so that other researchers can access it, but your name and other personal information will not be included. Usually, no one would be able to look at your genome and link it to you as a person. However, if another database exists that also has information on your genome and your name, someone might be able to compare the databases and identify you. If others found out, it could lead to discrimination or other problems. The risk of this is very small.

10. What are the risks of genetic testing?

It is unlikely, but the genetic tests done on your samples could show you may be at risk for certain diseases. If others found out, it could lead to discrimination or other problems. However, it is almost impossible for you or others to know your test results from the genetic testing. The results are not part of your study records and are not given to you.

11. Who will have access to my information in studies using my extra samples?

People who may see your information are:

- Researchers who use your extra samples and information for other research
- Government agencies that fund or monitor the research using your extra samples and information
- Any regulatory agency that reviews clinical trials
- The researcher's Institutional Review Board or Ethics Committee
- The people who work with the researcher

All of these people will do their best to protect your information. The results of any new studies that use your extra samples and information may be published. No publication will use your name or identify you personally.

Questions

12. If you have questions or problems about allowing your samples and information to be used in other studies, use the following important contacts.

If you have questions about the use of your samples or information or if you want to change your mind about their use, contact
[name or title and telephone number of the investigator or other study staff].

If you think you may have been harmed because of studies using your samples or information, contact
[name or title and telephone number of the investigator or other study staff].

If you have questions about your rights as a research participant, contact
[name or title and telephone number of person on IRB/EC .

Paragraph for South African sites. The study has been structured in accordance with the Declaration of Helsinki (last updated October 2013) which deals with the recommendations guiding doctors in biomedical research involving human participants, the Ethics in Health Research: Principles, Structures and Processes Second Edition 2015, and Guidelines for Good Practice in the Conduct of Clinical Trials in Human Participants in South Africa. We can provide you with copies of these guidelines if you wish to review them.

Remainder of section for South African sites only.

You can reach a study staff member 24 hours a day at [telephone number].

If you have questions about this trial you should first discuss them with your doctor or the ethics committee (contact details as provided on this form). After you have consulted your doctor or the ethics committee and if they have not provided you with answers to your satisfaction, you should write to the South African Healthcare Products Regulatory Authority (SAHPRA) at:

The Acting Chief Executive Officer
South African Health Products Regulatory Authority
Private Bag X828
PRETORIA
0001

Tel: (012) 842 7582/3

e-mail: portia.nkambule@sahpra.org.za

13. Please choose only one of the options below and write your initials or make your mark in the box next to it. Whatever you choose, the HVTN and HPTN keep track of your decision about how your samples and information can be used. You can change your mind after signing this form.

☐

I allow my extra samples and information to be used for other studies related to HIV, vaccines, monoclonal antibodies, the immune system, and other diseases. This may include genetic testing and keeping my cells growing over time.

OR

☐

I agree to the option above *and* also to allow my extra samples and information to be used in genome wide studies.

OR

☐

I do not allow my extra samples to be used in any other studies. This includes not allowing genetic testing, growing more of my cells, or genome wide studies.

Participant's name (print)	Participant's signature or mark	Date	Time
Clinic staff conducting consent discussion (print)	Clinic staff signature	Date	Time

For participants who are unable to read or write, a witness should complete the signature block below:

Witness's name (print)	Witness's signature	Date	Time
------------------------	---------------------	------	------

*Witness is impartial and was present for the entire discussion of this consent form.

Appendix D Tables of procedures for sample informed consent form

Table of procedures for Part 1: Screening and interrupting your HIV treatment

			Time after interrupting HIV treatment																		
	Screening	HIV treatment switch ^a	Pre-interruption visit about 4 weeks later	Day 0	Week 1	Week 2	Week 3	~1 month	Week 5	Week 6	Week 7	~2 months	Week 10	~3 months	Week 14	~4 months	Week 18	~5 months	Week 22		
Study procedures																					
HIV treatment switch (if required)		✓	✓																		
Medical history	✓																				
Complete physical exam	✓			✓																	
Brief physical exam		✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Pregnancy test and contraception review ^b	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Transmission risk reduction counseling	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Interview/questionnaire	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
STI testing (blood, urine, cervical/vaginal swabs) ^c	✓			✓				✓				✓		✓		✓		✓			
TB test	✓																				
Blood drawn	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		

Procedures in gray only for participants switching HIV treatments.

^a We will contact you about 2 weeks after you start the new HIV treatment to check to see if you have had any side effects or have other concerns.

^b Participants who have reached menopause or who had a hysterectomy, oophorectomy, or tubal ligation verified by medical records are not required to have a pregnancy test or contraception review.

^c In addition to STI testing at the checked visits, we will test at other visits if you show symptoms of an STI.

^d Extra visits every 3 months for people do not meet criteria for moving to Part 2 or Part 3.

Table of procedures for Part 1: Screening and interrupting your HIV treatment (continued)

	Time after interrupting HIV treatment								
	~6 months	Week 28	Week 32	Week 36	~ 9 months	Week 44	Week 48	1 year	Extra visits ^d
Study procedures									
HIV treatment switch (if required)									
Medical history									
Complete physical exam									
Brief physical exam	✓	✓	✓	✓	✓	✓	✓	✓	✓
Pregnancy test and contraception review ^b	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling	✓	✓	✓	✓	✓	✓	✓	✓	✓
Interview/questionnaire	✓	✓	✓	✓	✓	✓	✓	✓	✓
STI testing (blood, urine, cervical/vaginal swabs) ^c	✓	✓	✓	✓	✓	✓	✓	✓	✓
TB test									
Blood Drawn	✓	✓	✓	✓	✓	✓	✓	✓	✓

^a We will contact you about 2 weeks after you start the new HIV treatment to check to see if you have had any side effects or have other concerns.

^b Participants who have reached menopause or who had a hysterectomy, oophorectomy, or tubal ligation verified by medical records are not required to have a pregnancy test or contraception review.

^c In addition to STI testing at the checked visits, we will test at other visits if you show symptoms of an STI.

^d Extra visits every 3 months for people do not meet criteria for moving to Part 2 or Part 3.

Table of procedures for Part 2: Monitoring your health and your HIV

	Time after starting Part 2															
	Day 0	Week 1	Week 2	Week 3	~1 month	Week 5	Week 6	Week 7	~2 months	Week 10	~3 months	Week 14	~4 months	Week 18	~5 months	Week 22
Study procedures																
Complete physical exam	✓															
Brief physical exam		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Pregnancy test and contraception review ^a	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Interview/questionnaire	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
STI testing (blood, urine, cervical/vaginal swabs) ^b	✓				✓				✓		✓		✓		✓	
Blood drawn	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

^a Participants who have reached menopause or who had a hysterectomy, oophorectomy, or tubal ligation verified by medical records are not required to have a pregnancy test or contraception review.

^b In addition to STI testing at the checked visits, we will test at other visits if you show symptoms of an STI.

^c Extra visits every 3 months for people do not meet criteria for moving to Part 3.

Table of procedures for Part 2: Monitoring your health and your HIV (continued)

	Time after starting Part 2										
	Week 24	6 months	Week 28	Week 30	Week 32	Week 34	Week 36	~9 months	Week 44	Week 48	1 year
Study procedures											
Complete physical exam											
Brief physical exam	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Pregnancy test and contraception review ^a	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Interview/questionnaire	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
STI testing (blood, urine, cervical/vaginal swabs) ^b	✓		✓		✓		✓	✓	✓	✓	✓
Blood drawn	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

^a Participants who have reached menopause or who had a hysterectomy, oophorectomy, or tubal ligation verified by medical records are not required to have a pregnancy test or contraception review.

^b In addition to STI testing at the checked visits, we will test if you show symptoms of an STI.

^c Extra visits every 3 months for people do not meet criteria for moving to Part 3.

Table of procedures for Part 3: Restart HIV treatment

	Time after restarting HIV treatment												
	Week 0	Week 2	~1 month	Week 6	~2 months	Week 10	Week 12	~4 months	Week 20	~6 months	Week 28	~9 months	1 year
Study procedures													
Complete physical exam													✓
Brief physical exam	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Pregnancy test & contraception review ^a	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling	✓	✓	✓		✓		✓		✓		✓		✓
Interview/Questionnaire	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
STI testing (blood, urine, cervical/vaginal swabs) ^b	✓		✓		✓		✓	✓ ^c	✓ ^c	✓	✓ ^c	✓	
Blood Drawn	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

^a Participants who have reached menopause or who had a hysterectomy, oophorectomy, or tubal ligation verified by medical records are not required to have a pregnancy test or contraception review. Pregnancy test and contraceptive review are not required once viral load drops to undetectable after restarting HIV treatment.

^b In addition to STI testing at the checked visits, we will test if you show symptoms of an STI.

^c STI testing is not required at this visit if viral load has returned to undetectable.

Appendix E Laboratory procedures—Schedule 1: Monitoring ATI

Procedure	Ship to ¹	Assay location ^{1,2}	Tube Type ³	Tube size (vol. capacity) ³	Visit: Days on ATI: Weeks on ATI:	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
					Screening visit ¹⁵	ART Switch ⁸	ATI qualification ⁹	D0 W0	D7 W1	D14 W2	D21 W3	D28 W4	D35 W5	D42 W6	D49 W7	D56 W8	D70 W10	D84 W12	D98 W14	D112 W16	D126 W18	D140 W20	D154 W22		
								ATI																	
BLOOD COLLECTION																									
Screening or diagnostic assays																									
HIV PCR viral load ¹³	Local labs	Local labs	EDTA	6mL	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	
CD4+/CD8+ T-cell count ¹⁴	Local labs	Local labs	EDTA	4mL	4	4	4	4	—	4	—	4	—	4	—	4	—	4	—	4	—	4	—	4	—
HBsAg/anti-HCV ⁴	Local labs	Local labs	SST	5mL	10	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
QuantIFERON TB testing ⁵	Local labs	Local labs	QFT Gold/Gold-Plus	1mL	4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Safety labs																									
Hgb / ANC / PLT	Local labs	Local labs	EDTA	4mL	4	4	4	4	—	—	—	—	4	—	—	—	4	—	4	—	4	—	4	—	
ALT / direct bilirubin / eGFR	Local labs	Local labs	SST	5mL	5	5	5	5	—	—	—	—	5	—	—	—	5	—	5	—	5	—	5	—	
Syphilis ¹⁰	Local labs	Local labs	SST	5mL	5	—	—	5	—	—	—	—	5	—	—	—	5	—	5	—	5	—	5	—	
Hormone Levels																									
Hormone panel	Local labs	Local labs	SST	5mL	—	—	—	w	—	—	—	w	—	—	—	—	w	—	w	—	w	—	—	—	
Drug levels/detection																									
ARV detection by dry blood spot	CSR	HVTN labs	EDTA	2mL	—	—	—	—	—	—	—	2	—	—	—	—	2	—	2	—	2	—	2	—	
Immunogenicity & Virologic Assays																									
Cellular assays																									
ICS	CSR	HVTN labs	ACD	8.5mL	—	—	—	42.5	—	—	—	—	42.5	—	—	—	42.5	—	42.5	—	42.5	—	42.5	—	
Phenotyping	CSR	HVTN labs	ACD	8.5mL	—	—	—	z	—	—	—	—	z	—	—	—	z	—	z	—	z	—	z	—	
Humoral assays																									
Neutralizing antibody	CSR	HVTN labs	SST	8.5mL	—	—	—	8.5	—	—	—	—	8.5	—	—	—	8.5	—	8.5	—	8.5	—	8.5	—	
FcR-mediated effector functions ⁶	CSR	HVTN labs	SST	8.5mL	—	—	—	y	—	—	—	—	y	—	—	—	y	—	y	—	y	—	y	—	
HIV reservoir assays	CSR	TBD	ACD	8.5mL	—	—	—	51	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Storage																									
Serum	CSR		SST	8.5mL	—	—	—	8.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Plasma	CSR		EDTA	10mL	—	—	—	—	—	—	—	—	—	—	—	—	10	—	10	—	10	—	10	—	
PBMC	CSR		ACD	8.5mL	—	—	—	—	—	—	—	—	—	—	—	—	17	—	17	—	17	—	17	—	
Visit total					38	19	19	134.5	6	10	6	77	6	10	6	104	6	104	6	104	6	104	6	6	
56-Day total					38	57	76	210.5	216.5	226.5	232.5	309.5	315.5	325.5	331.5	397.5	225	313	236	324	226	324	226	226	
URINE COLLECTION																									
Pregnancy Test ¹⁶	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Chlamydia/gonorrhea ⁷	Local labs	Local labs			X	—	—	X	—	—	—	X	—	—	—	X	—	X	—	X	—	X	—	—	
CERVICAL/VAGINAL SWAB COLLECTION																									
Chlamydia/gonorrhea ⁷	Local labs	Local labs			X	—	—	X	—	—	—	X	—	—	—	X	—	X	—	X	—	X	—	—	
Trichomonas vaginalis ⁷	Local labs	Local labs			X	—	—	X	—	—	—	X	—	—	—	X	—	X	—	X	—	X	—	—	

¹ CSR = central specimen repository.

² HVTN Laboratories include: Fred Hutchinson Cancer Research Center (Seattle, Washington, USA); Duke University Medical Center (Durham, North Carolina, USA); South African Immunology Laboratory-National Institute for Communicable Diseases (SAIL-NICD, Johannesburg, South Africa); University of Cape Town (Cape Town, South Africa); Cape Town HVTN Immunology Laboratory (CHIL, Cape Town, South Africa).

Non-HVTN laboratories: TBD.

³ Local labs may assign appropriate alternative tube types for locally performed tests.

⁴ HCV RNA PCR testing will be performed as a reflex test if indicated by anti-HCV antibody results and may require an additional blood collection.

⁵ Tuberculin skin test (TST) will be performed if QuantiFERON TB testing is not available. See Procedures at CRS ([Appendix H](#)).

⁶ FcR-mediated effector function assays may include ADCC, virion capture, and phagocytosis assays.

⁷ Chlamydia/gonorrhea testing will be done on EITHER urine OR a cervical/vaginal swab; Trichomonas testing will be done on cervical/vaginal swab. In addition to STI testing at the marked visits, STI testing may occur at any visit if clinically indicated. In addition to the listed specimen types (ie., cervical/vaginal swabs), chlamydia/gonorrhea testing may occur on rectal swabs if clinically indicated (see HVTN 805/HPTN 093 SSP for details).

⁸ The "ART switch" phase will only be performed for participants on NNRTIs. These participants will be considered enrolled on the first day of the new ART medication.

⁹ The ATI Qualification visit specimens must be obtained at least 28 days after ART switch. If needed, VL retesting may continue until viral suppression has been achieved (up to 84 days after ART switch). The last ATI qualification procedures must take place no more than 14 days prior to visit 4 (see HVTN 805/HPTN 093 SSP for more information).

¹⁰ In addition to syphilis testing at the marked visits, syphilis testing may occur at any visit if clinically indicated.

- ¹¹ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 27 continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met criteria to transition to Schedule 2 or Schedule 3 (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹² Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 27 continuing up to 3 years of this schedule, and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met criteria to transition to Schedule 2 or Schedule 3 (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹³ A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first VL result ≥ 200 copies/mL (see Protocol Section 3.3.1 and HVTN 805/HPTN 093 SSP for details).
- ¹⁴ A confirmatory sample should be at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see Protocol Section 3.3.2 and HVTN 805/HPTN 093 SSP for details).
- ¹⁵ Screening visit specimens for participants not undergoing an NNRTI switch should be obtained no later than 2 weeks before Visit 4 (see HVTN 805/HPTN 093 SSP for more information).
- ¹⁶ For persons capable of becoming pregnant, pregnancy test may be performed on urine or blood specimens.
- ¹⁷ At an early termination visit for a withdrawn or terminated participant (see Protocol Section 6.5), blood should be drawn as shown for Extended follow-up visit type A (see HVTN 805/HPTN 093 SSP for more information).
- w = SST blood collected for syphilis testing will also cover specimen needs for HBsAg and anti-HCV screening testing. Hormone panel is defined in Protocol Sections 6.1 (Schedule 1: Monitoring ATI) and 6.2 (Schedule 2: Monitoring ATI with viremia).
- y = SST blood collected for neutralizing antibody will also cover specimen needs for FcR-mediated effector functions; no separate blood draw is needed.
- z = PBMC blood collected for ICS will also cover specimen needs for phenotyping; no separate blood draw is needed.

Laboratory procedures—Schedule 1: Monitoring ATI (continued)

Procedure	Ship to ¹	Assay location ²	Tube Type ³	Tube size (vol. capacity) ³	Visit:	20	21	22	23	24	25	26	27	Visit Type A ^{11,17}	Visit Type B ¹²	Total
					Days on ATI:	D168	D196	D224	D252	D280	D308	D336	D364			
					Weeks on ATI:	W24	W28	W32	W36	W40	W44	W48	W52			
BLOOD COLLECTION																
Screening or diagnostic assays																
HIV PCR viral load ¹³	Local labs	Local labs	EDTA	6mL	6	6	6	6	6	6	6	6	6	6	6	174
CD4+/CD8+ T-cell count ¹⁴	Local labs	Local labs	EDTA	4mL	4	4	4	4	—	4	—	4	4	4	4	76
HBsAg/anti-HCV	Local labs	Local labs	SST	5mL	—	—	—	—	—	—	—	—	—	—	—	10
QuantiFERON TB testing ⁵	Local labs	Local labs	QFT Gold/Gold-Plus	1mL	—	—	—	—	—	—	—	—	—	—	—	4
Safety labs																
Hgb / ANC / PLT	Local labs	Local labs	EDTA	4mL	4	4	4	4	—	4	—	4	4	4	4	68
ALT / direct bilirubin / eGFR	Local labs	Local labs	SST	5mL	5	5	5	5	—	5	—	5	5	5	5	85
Syphilis ¹⁰	Local labs	Local labs	SST	5mL	5	5	5	5	5	5	5	5	5	5	5	85
Hormone Levels																
Hormone panel	Local labs	Local labs	SST	5mL	—	—	—	w	—	—	—	—	—	—	—	0
Drug levels/detection																
ARV detection by dry blood spot	CSR	HVTN labs	EDTA	2mL	2	2	2	2	2	2	2	2	2	2	2	30
Immunogenicity & Virologic Assays																
Cellular assays																
ICS	CSR	HVTN labs	ACD	8.5mL	42.5	42.5	42.5	42.5	42.5	42.5	42.5	42.5	42.5	—	—	595
Phenotyping	CSR	HVTN labs	ACD	8.5mL	z	z	z	z	z	z	z	z	z	—	—	0
Humoral assays																
Neutralizing antibody	CSR	HVTN labs	SST	8.5mL	8.5	8.5	8.5	8.5	8.5	8.5	8.5	8.5	8.5	—	—	119
FcR-mediated effector functions ⁶	CSR	HVTN labs	SST	8.5mL	y	y	y	y	y	y	y	y	y	—	—	0
HIV reservoir assays	CSR	TBD	ACD	8.5mL	—	—	—	—	—	—	—	—	—	—	—	51
Storage																
Serum	CSR		SST	8.5mL	—	—	—	—	—	—	—	—	—	8.5	—	17
Plasma	CSR		EDTA	10mL	10	10	10	10	10	10	10	10	10	—	—	130
PBMC	CSR		ACD	8.5mL	17	17	17	17	17	17	17	17	17	59.5	—	263.5
Visit total					104	104	104	104	91	104	91	104	104	104	26	
56-Day total					324	318	312	312	299	299	286	299	104	26		
URINE COLLECTION																
Pregnancy Test ¹⁶	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	
Chlamydia/gonorrhea ⁷	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	
CERVICAL/VAGINAL SWAB COLLECTION																
Chlamydia/gonorrhea ⁷	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	
Trichomonas vaginalis ⁷	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	

¹ CSR = central specimen repository.

² HVTN Laboratories include: Fred Hutchinson Cancer Research Center (Seattle, Washington, USA); Duke University Medical Center (Durham, North Carolina, USA); South African Immunology Laboratory-National Institute for Communicable Diseases (SAIL-NICD, Johannesburg, South Africa); University of Cape Town (Cape Town, South Africa); Cape Town HVTN Immunology Laboratory (CHIL, Cape Town, South Africa).

Non-HVTN laboratories: TBD.

³ Local labs may assign appropriate alternative tube types for locally performed tests.

⁴ HCV RNA PCR testing will be performed as a reflex test if indicated by anti-HCV antibody results and may require an additional blood collection..

⁵ Tuberculin skin test (TST) will be performed if QuantiFERON TB testing is not available. See Procedures at CRS (Appendix H).

⁶ FcR-mediated effector function assays may include ADCC, virion capture, and phagocytosis assays.

⁷ Chlamydia/gonorrhea testing will be done on EITHER urine OR a cervical/vaginal swab; Trichomonas testing will be done on cervical/vaginal swab. In addition to STI testing at the marked visits, STI testing may occur at any visit if clinically indicated. In addition to the listed specimen types (ie., cervical/vaginal swabs), chlamydia/gonorrhea testing may occur on rectal swabs if clinically indicated (see HVTN 805/HPTN 093 SSP for details).

⁸ The "ART switch" phase will only be performed for participants on NNRTIs. These participants will be considered enrolled on the first day of the new ART medication.

⁹ The ATI Qualification visit specimens must be obtained at least 28 days after ART switch. If needed, VL retesting may continue until viral suppression has been achieved (up to 84 days after ART switch). The last ATI qualification procedures must take place no more than 14 days prior to visit 4 (see HVTN 805/HPTN 093 SSP for more information).

¹⁰ In addition to syphilis testing at the marked visits, syphilis testing may occur at any visit if clinically indicated.

- ¹¹ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 27 continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met criteria to transition to Schedule 2 or Schedule 3 (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹² Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 27 continuing up to 3 years of this schedule, and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met criteria to transition to Schedule 2 or Schedule 3 (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹³ A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first VL result ≥ 200 copies/mL (see Protocol Section 3.3.1 and HVTN 805/HPTN 093 SSP for details).
- ¹⁴ A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see Protocol Section 3.3.2 and HVTN 805/HPTN 093 SSP for details).
- ¹⁵ Screening visit specimens for participants not undergoing an NNRTI switch should be obtained no later than 2 weeks before Visit 4 (see HVTN 805/HPTN 093 SSP for more information).
- ¹⁶ For persons capable of becoming pregnant, pregnancy test may be performed on urine or blood specimens.
- ¹⁷ At an early termination visit for a withdrawn or terminated participant (see Protocol Section 6.5), blood should be drawn as shown for Extended follow-up visit type A (see HVTN 805/HPTN 093 SSP for more information).
- w = SST blood collected for syphilis testing will also cover specimen needs for HBsAg and anti-HCV screening testing. Hormone panel is defined in Protocol Sections 6.1 (Schedule 1: Monitoring ATI) and 6.2 (Schedule 2: Monitoring ATI with viremia).
- y = SST blood collected for neutralizing antibody will also cover specimen needs for FcR-mediated effector functions; no separate blood draw is needed.
- z = PBMC blood collected for ICS will also cover specimen needs for phenotyping; no separate blood draw is needed.

Appendix F Laboratory procedures—Schedule 2: Monitoring ATI with viremia

Procedure	Ship to ¹	Assay location ^{1,2}	Tube Type ³	Tube size (vol. capacity) ³	Visit:	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55
					Days on ATI with viremia	D0	D7	D14	D21	D28	D35	D42	D49	D56	D70	D84	D98	D112	D126	D140	D154
					Weeks on ATI with viremia	W0	W1	W2	W3	W4	W5	W6	W7	W8	W10	W12	W14	W16	W18	W20	W22
BLOOD COLLECTION																					
Screening or diagnostic assays																					
HIV PCR viral load ¹⁰	Local labs	Local labs	EDTA	6mL	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
CD4+/CD8+ T-cell count ¹¹	Local labs	Local labs	EDTA	4mL	4	—	4	—	4	—	4	—	4	—	4	—	4	—	4	—	4
Safety labs																					
Hgb / ANC / PLT	Local labs	Local labs	EDTA	4mL	4	—	—	—	—	4	—	—	—	4	—	4	—	4	—	4	—
ALT / direct bilirubin / eGFR	Local labs	Local labs	SST	5mL	5	—	—	—	—	5	—	—	—	5	—	5	—	5	—	5	—
Syphilis ⁷	Local labs	Local labs	SST	5mL	5	—	—	—	—	5	—	—	—	5	—	5	—	5	—	5	—
Hormone Levels																					
Hormone panel	Local labs	Local labs	SST	5mL	w	—	—	—	—	w	—	—	—	w	—	w	—	w	—	—	—
Drug levels/detection																					
ARV detection by dry blood spot	CSR	HVTN labs	EDTA	2mL	—	—	—	—	—	2	—	—	—	2	—	2	—	2	—	2	—
Immunogenicity & Virologic Assays																					
Cellular assays																					
ICS	CSR	HVTN labs	ACD	8.5mL	42.5	—	—	42.5	—	42.5	—	—	—	42.5	—	42.5	—	42.5	—	42.5	—
Phenotyping	CSR	HVTN labs	ACD	8.5mL	z	—	—	z	—	z	—	—	—	z	—	z	—	z	—	z	—
Humoral assays																					
Neutralizing antibody	CSR	HVTN labs	SST	8.5mL	8.5	—	—	8.5	—	8.5	—	—	—	8.5	—	8.5	—	8.5	—	8.5	—
FcR-mediated effector functions ⁴	CSR	HVTN labs	SST	8.5mL	y	—	—	y	—	y	—	—	—	y	—	y	—	y	—	y	—
Storage																					
Serum	CSR		SST	8.5mL	8.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Plasma	CSR		EDTA	10mL	10	—	—	—	—	—	—	—	—	10	—	10	—	10	—	10	—
PBMC	CSR		ACD	8.5mL	—	—	—	—	—	—	—	—	—	17	—	17	—	17	—	17	—
Visit total					93.5	6	61	6	77	6	10	6	104	6	104	6	104	6	104	6	104
56-Day total^{5,10,11}					93.5	99.5	160.5	166.5	243.5	249.5	259.5	265.5	369.5	276	313	236	324	226	324	226	226
URINE COLLECTION																					
Pregnancy Test ¹²	Local labs	Local labs			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Chlamydia/gonorrhea ⁵	Local labs	Local labs			X	—	—	—	X	—	—	—	—	X	—	X	—	X	—	X	—
CERVICAL/VAGINAL SWAB COLLECTION																					
Chlamydia/gonorrhea ⁵	Local labs	Local labs			X	—	—	—	X	—	—	—	—	X	—	X	—	X	—	X	—
Trichomonas vaginalis ⁵	Local labs	Local labs			X	—	—	—	X	—	—	—	—	X	—	X	—	X	—	X	—

¹ CSR = central specimen repository.

² HVTN Laboratories include: Fred Hutchinson Cancer Research Center (Seattle, Washington, USA); Duke University Medical Center (Durham, North Carolina, USA); South African Immunology Laboratory-National Institute for Communicable Diseases (SAIL-NICD, Johannesburg, South Africa); University of Cape Town (Cape Town, South Africa); Cape Town HVTN Immunology Laboratory (CHIL, Cape Town, South Africa).

Non-HVTN laboratories: TBD.

³ Local labs may assign appropriate alternative tube types for locally performed tests.

⁴ FcR-mediated effector function assays may include ADCC, virion capture, and phagocytosis assays.

⁵ Chlamydia/gonorrhea testing will be done on EITHER urine OR a cervical/vaginal swab; Trichomonas testing will be done on cervical/vaginal swab. In addition to STI testing at the marked visits, STI testing may occur at any visit if clinically indicated. In addition to the listed specimen types (ie., cervical/vaginal swabs), chlamydia/gonorrhea testing may occur on rectal swabs if clinically indicated (see HVTN 805/HPTN 093 SSP for details).

⁶ The 56-day blood draw limit does not include visits from Schedule 1; please see HVTN 805/HPTN 093 SSP for details on prioritizing collections to avoid exceeding the 56-day limit.

⁷ In addition to syphilis testing at the marked visits, syphilis testing may occur at any visit if clinically indicated.

⁸ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 66, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁹ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 66, continuing up to 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

¹⁰ Additional weekly viral load monitoring may be required between weeks 8 and 24; after week 24, a confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first VL result ≥ 200 copies/mL (see Protocol Section 3.3.3 and HVTN 805/HPTN 093 SSP for details). The 56-day blood draw limit does not include up to 10mL blood collected per visit for this additional monitoring; however, the 56-day limit is not exceeded at any visit by these collections.

¹¹ A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see Protocol Section 3.3.3 and HVTN 805/HPTN 093 SSP for details).

¹² For persons capable of becoming pregnant, pregnancy test may be performed on urine or blood specimens.

¹³ At an early termination visit for a withdrawn or terminated participant (see Protocol Section 6.5), blood should be drawn as shown for Extended follow-up visit type A (see HVTN 805/HPTN 093 SSP for more information).

w = SST blood collected for syphilis testing will also cover specimen needs for HBsAg and anti-HCV screening testing. Hormone panel is defined in Protocol Sections 6.1 (Schedule 1: Monitoring ATI) and 6.2 (Schedule 2: Monitoring ATI with viremia).

y = SST blood collected for neutralizing antibody will also cover specimen needs for FcR-mediated effector functions; no separate blood draw is needed.

z = PBMC blood collected for ICS will also cover specimen needs for phenotyping; no separate blood draw is needed.

Laboratory procedures—Schedule 2: Monitoring ATI with viremia (continued)

Procedure	Ship to ¹	Assay location ^{1,2}	Tube Type ³	Tube size (vol. capacity) ³	Visit: Days on ATI with viremia	56	57	58	59	60	61	62	63	64	65	66	Visit Type A ^{8,13}	Visit type B ⁹	Total
					Weeks on ATI with viremia	D168 W24	D182 W26	D196 W28	D210 W30	D224 W32	D238 W34	D252 W36	D280 W40	D308 W44	D336 W48	D364 W52			
BLOOD COLLECTION																			
Screening or diagnostic assays																			
HIV PCR viral load ¹⁰	Local labs	Local labs	EDTA	6mL	6	6	6	6	6	6	6	6	6	6	6	6	6	6	174
CD4+/CD8+ T-cell count ¹¹	Local labs	Local labs	EDTA	4mL	4	—	4	—	4	—	4	—	4	—	4	—	4	4	64
Safety labs																			
Hgb / ANC / PLT	Local labs	Local labs	EDTA	4mL	4	—	4	—	4	—	4	—	4	—	4	—	4	4	56
ALT / direct bilirubin / eGFR	Local labs	Local labs	SST	5mL	5	—	5	—	5	—	5	—	5	—	5	—	5	5	70
Syphilis ⁷	Local labs	Local labs	SST	5mL	5	—	5	—	5	—	5	—	5	—	5	—	5	5	80
Hormone Levels																			
Hormone panel	Local labs	Local labs	SST	5mL	—	—	—	—	—	—	w	—	—	—	—	—	—	—	0
Drug levels/detection																			
ARV detection by dry blood spot	CSR	HVTN labs	EDTA	2mL	2	—	2	—	2	—	2	2	2	2	2	2	2	2	30
Immunogenicity & Virologic Assays																			
Cellular assays																			
ICS	CSR	HVTN labs	ACD	8.5mL	42.5	—	42.5	—	42.5	—	42.5	42.5	42.5	42.5	42.5	42.5	—	—	637.5
Phenotyping	CSR	HVTN labs	ACD	8.5mL	z	—	z	—	z	—	z	z	z	z	z	z	—	—	0
Humoral assays																			
Neutralizing antibody	CSR	HVTN labs	SST	8.5mL	8.5	—	8.5	—	8.5	—	8.5	8.5	8.5	8.5	8.5	8.5	—	—	127.5
FcR-mediated effector functions ⁴	CSR	HVTN labs	SST	8.5mL	y	—	y	—	y	—	y	y	y	y	y	y	—	—	0
Storage																			
Serum	CSR		SST	8.5mL	—	—	—	—	—	—	—	—	—	—	—	—	8.5	—	17
Plasma	CSR		EDTA	10mL	10	—	10	—	10	—	10	10	10	10	10	10	10	—	140
PBMC	CSR		ACD	8.5mL	17	—	17	—	17	—	17	17	17	17	17	17	59.5	—	263.5
Visit total						104	6	104	6	104	6	104	91	104	91	104	104	26	
56-Day total^{6,10,11}						324	226	324	226	324	226	324	305	299	286	299	104	26	
URINE COLLECTION																			
Pregnancy Test ¹²	Local labs	Local labs				X	X	X	X	X	X	X	X	X	X	X	X	X	
Chlamydia/gonorrhea ⁵	Local labs	Local labs				X	—	X	—	X	—	X	X	X	X	X	X	X	
CERVICAL/VAGINAL SWAB COLLECTION																			
Chlamydia/gonorrhea ⁵	Local labs	Local labs				X	—	X	—	X	—	X	X	X	X	X	X	X	
Trichomonas vaginalis ⁵	Local labs	Local labs				X	—	X	—	X	—	X	X	X	X	X	X	X	

¹ CSR = central specimen repository.

² HVTN Laboratories include: Fred Hutchinson Cancer Research Center (Seattle, Washington, USA); Duke University Medical Center (Durham, North Carolina, USA); South African Immunology Laboratory-National Institute for Communicable Diseases (SAIL-NICD, Johannesburg, South Africa); University of Cape Town (Cape Town, South Africa); Cape Town HVTN Immunology Laboratory (CHIL, Cape Town, South Africa).

Non-HVTN laboratories: TBD.

³ Local labs may assign appropriate alternative tube types for locally performed tests.

⁴ FcR-mediated effector function assays may include ADCC, virion capture, and phagocytosis assays.

⁵ Chlamydia/gonorrhea testing will be done on EITHER urine OR a cervical/vaginal swab; Trichomonas testing will be done on cervical/vaginal swab. In addition to STI testing at the marked visits, STI testing may occur at any visit if clinically indicated. In addition to the listed specimen types (ie., cervical/vaginal swabs), chlamydia/gonorrhea testing may occur on rectal swabs if clinically indicated (see HVTN 805/HPTN 093 SSP for details).

⁶ The 56-day blood draw limit does not include visits from Schedule 1; please see HVTN 805/HPTN 093 SSP for details on prioritizing collections to avoid exceeding the 56-day limit.

⁷ In addition to syphilis testing at the marked visits, syphilis testing may occur at any visit if clinically indicated.

- ⁸ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 66, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ⁹ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 66, continuing up to 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹⁰ Additional weekly viral load monitoring may be required between weeks 8 and 24; after week 24, a confirmatory sample should be drawn as soon as possible following the first VL result ≥ 200 copies/mL (see Protocol Section 3.3.3 for details). The 56-day blood draw limit does not include up to 10mL blood collected per visit for this additional monitoring; however, the 56-day limit is not exceeded at any visit by these collections.
- ¹¹ A confirmatory sample should be drawn at the next visit (within approximately 1-2 weeks) following the first CD4+ T-cell count < 350 cells/mm³ (see Protocol Section 3.3.3 and HVTN 805/HPTN 093 SSP for details).
- ¹² For persons capable of becoming pregnant, pregnancy test may be performed on urine or blood specimens.
- ¹³ At an early termination visit for a withdrawn or terminated participant (see Protocol Section 6.5), blood should be drawn as shown for Extended follow-up visit type A (see HVTN 805/HPTN 093 SSP for more information).
- w = SST blood collected for syphilis testing will also cover specimen needs for HBsAg and anti-HCV screening testing. Hormone panel is defined in Protocol Sections 6.1 (Schedule 1: Monitoring ATI) and 6.2 (Schedule 2: Monitoring ATI with viremia).
- y = SST blood collected for neutralizing antibody will also cover specimen needs for FcR-mediated effector functions; no separate blood draw is needed.
- z = PBMC blood collected for ICS will also cover specimen needs for phenotyping; no separate blood draw is needed.

Appendix G Laboratory procedures—Schedule 3: Follow-up on ART

Procedure	Ship to ¹	Assay location ^{1,2}	Tube Type ³	Tube size (vol. capacity) ³	Visit:	80 ⁷	81	82	83	84	85	86	87	88	89	90	91 ¹²	92	Total
					Days post ART re-initiation	D0	D14	D28	D42	D56	D70	D84	D112	D140	D168	D196	D280	D364	
					Weeks post ART re-initiation:	W0	W2	W4	W6	W8	W10	W12	W16	W20	W24	W28	W40	W52	
					ART re-initiation visit														
BLOOD COLLECTION																			
Screening or diagnostic assays																			
HIV PCR viral load	Local labs	Local labs	EDTA	6mL	6	6	6	6	6	6	6	6	6	6	6	6	6	6	78
CD4+/CD8+ T-cell count	Local labs	Local labs	EDTA	4mL	4	—	4	—	4	—	4	—	4	—	4	4	4	4	32
Safety labs																			
Hgb / ANC / PLT	Local labs	Local labs	EDTA	4mL	—	—	4	—	4	—	4	—	—	4	—	4	—	—	20
ALT / direct bilirubin / eGFR	Local labs	Local labs	SST	5mL	—	—	5	—	5	—	5	—	—	5	—	5	—	—	25
Syphilis ⁹	Local labs	Local labs	SST	5mL	5	—	5	—	5	—	5	5 ¹³	5 ¹³	5	5 ¹³	5	—	—	45
HIV genotypic antiretroviral resistance ⁸	CSR	HVTN Labs	EDTA	6mL	—	—	—	—	12	—	—	—	—	—	—	—	—	—	12
Immunogenicity & Virologic Assays																			
Viral isolation/sequencing	CSR	HVTN Labs	EDTA	10mL	10	10	—	—	—	—	10	—	—	10	—	10	10	60	
Cellular assays																			
ICS	CSR	HVTN Labs	ACD	8.5mL	—	—	—	—	—	—	42.5	—	—	42.5	—	42.5	42.5	170	
Phenotyping	CSR	HVTN Labs	ACD	8.5mL	—	—	—	—	—	—	z	—	—	z	—	z	z	0	
Humoral assays																			
Neutralizing antibody	CSR	HVTN Labs	SST	8.5mL	—	—	—	—	—	—	8.5	—	—	8.5	—	8.5	8.5	34	
FcR-mediated effector functions ⁴	CSR	HVTN Labs	SST	8.5mL	—	—	—	—	—	—	y	—	—	y	—	y	y	0	
HIV reservoir assays	CSR	TBD	ACD	8.5mL	34	—	—	—	—	—	—	—	—	51	—	51	51	187	
Storage																			
Serum	CSR		SST	8.5mL	8.5	—	—	—	—	—	8.5	—	—	8.5	—	8.5	8.5	42.5	
Plasma	CSR		EDTA	10mL	—	—	—	—	—	—	10	—	—	10	—	10	10	40	
PBMC	CSR		ACD	8.5mL	42.5	—	—	—	—	—	102	—	—	51	—	51	51	297.5	
Visit total					110	16	24	6	36	6	205.5	11	15	201.5	15	205.5	191.5		
56-Day total ⁵						126	150	156	192	88	277.5	258.5	231.5	227.5	231.5	205.5	191.5		
URINE COLLECTION																			
Pregnancy Test ¹¹	Local labs	Local labs			X	X ¹⁰	X ¹⁰	X ¹⁰	X ¹⁰	X ¹⁰	X	X ¹⁰	X ¹⁰	X	X ¹⁰	X	X		
Chlamydia/gonorrhea ⁶	Local labs	Local labs			X	—	X	—	X	—	X	X ¹³	X ¹³	X	X ¹³	X	—		
CERVICAL/VAGINAL SWAB COLLECTION																			
Chlamydia/gonorrhea ⁶	Local labs	Local labs			X	—	X	—	X	—	X	X ¹³	X ¹³	X	X ¹³	X	—		
Trichomonas vaginalis ⁵	Local labs	Local labs			X	—	X	—	X	—	X	X ¹³	X ¹³	X	X ¹³	X	—		

¹ CSR = central specimen repository.

² HVTN Laboratories include: Fred Hutchinson Cancer Research Center (Seattle, Washington, USA); Duke University Medical Center (Durham, North Carolina, USA); South African Immunology Laboratory-National Institute for Communicable Diseases (SAIL-NICD, Johannesburg, South Africa); University of Cape Town (Cape Town, South Africa); Cape Town HVTN Immunology Laboratory (CHIL, Cape Town, South Africa).

Non-HVTN laboratories: TBD.

³ Local labs may assign appropriate alternative tube types for locally performed tests.

⁴ FcR-mediated effector function assays may include ADCC, virion capture, and phagocytosis assays.

⁵ 56-day totals do not include visit totals from Schedule 1 or 2. See HVTN 805/HPTN 093 SSP for details on prioritizing collections to avoid exceeding the 56-day limit.

⁶ Chlamydia/gonorrhea testing will be done on EITHER urine OR a cervical/vaginal swab; Trichomonas testing will be done on cervical/vaginal swab. In addition to STI testing at the marked visits, STI testing may occur at any visit if clinically indicated. In addition to the listed specimen types (ie., cervical/vaginal swabs), chlamydia/gonorrhea testing may occur on rectal swabs if clinically indicated (see HVTN 805/HPTN 093 SSP for details).

⁷ Samples for visit 80 should be collected after ART re-initiation criteria have been met, but prior to ART re-initiation.

⁸ HIV antiretroviral resistance testing will be performed only if indicated by viral load results (see HVTN 805/HPTN 093 SSP).

⁹ In addition to syphilis testing at the marked visits, syphilis testing may occur at any visit if clinically indicated.

¹⁰ Pregnancy testing is not required if viral load has returned to undetectable.

¹¹ For persons capable of becoming pregnant, pregnancy test may be performed on urine or blood specimens.

¹² At an early termination visit for a withdrawn or terminated participant (see Protocol Section 6.5), blood should be drawn as shown for Visit 91 (see HVTN 805/HPTN 093 SSP for more information).

¹³ Syphilis, chlamydia and gonorrhea testing is not required at this visit if viral load has returned to undetectable.

y = SST blood collected for neutralizing antibody will also cover specimen needs for FcR-mediated effector functions; no separate blood draw is needed.

z = PBMC blood collected for ICS will also cover specimen needs for phenotyping; no separate blood draw is needed.

Appendix H Procedures at CRS—Schedule 1: Monitoring ATI

	Visit: Days on ATI: Weeks on ATI:	01 ¹ Screening	02 ART Switch ²	03 ATI qualification	04 ³ D0 W0	05 D7 W1	06 D14 W2	07 D21 W3	08 D28 W4	09 D35 W5	10 D42 W6	11 D49 W7	12 D56 W8	13 D70 W10	14 D84 W12	15 D98 W14	16 D112 W16	17 D126 W18	18 D140 W20	19 D154 W22
Study procedures																				
Screening consent (if used)		✓																		
Protocol consent		✓																		
Assessment of understanding		✓																		
Medical history		✓																		
Complete physical exam		✓			✓															
Targeted physical exam ⁴			✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Concomitant medications ⁴	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercurrent illness/adverse experience ⁴			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
ART re-initiation assessment ⁴						✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling ⁴	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Contraception status assessment ⁵	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Decision aid	✓																			
Decision-making assessment			✓		✓										✓					
Psychosocial assessment			✓		✓										✓					
Social impact assessment ⁴					✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Social impact assessment questionnaire															✓					
QuantiFERON tuberculosis test ⁶	✓																			
Confirm eligibility	✓																			
Specimen collection⁷	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ Screening may occur over the course of several contacts/visits up to and including the day of enrollment as defined in Section 6.1.2.

² For participants undergoing switch from NNRTI-based to protease- or integrase-based ART regimen. Participants undergoing an “ART switch” will be considered enrolled on first day of the new ART medication. The “ART switch” phase will not be performed for participants not on NNRTIs. For procedure timing during the “ART switch,” see Section 6.1.3 and HVTN 805/HPTN 093 SSP.

³ Enrollment visit for participants who do not undergo ART switch.

⁴ Procedure to be performed at interim visits held to draw confirmatory viral load samples or confirmatory samples for CD4+ T cell counts (see Sections 3.3.1, 3.3.2, and 3.3.3).

⁵ Contraception status assessment is required only for participants who can become pregnant (does not include those persons not of reproductive potential due to having undergone hysterectomy or bilateral oophorectomy [verified by medical records]).

⁶ If QuantiFERON TB testing cannot be performed, a tuberculin skin test (TST) should be conducted. Additional risk/clinical/diagnostic assessment may be performed at the discretion of the clinician to meet institutional standard of care for evaluation and treatment of latent TB.

⁷ For specimen collection requirements, see Appendix E.

⁸ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 27, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁹ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 27, continuing up 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section [3.3](#) and HVTN 805/HPTN 093 SSP for details).

¹⁰ At an early termination visit for a withdrawn or terminated participant, CRS staff should consider performing procedures specified for Extended follow-up visit Type A (see Section [6.5](#) and HVTN 805/HPTN 093 SSP for details).

Procedures at CRS—Schedule 1: Monitoring ATI (continued)

Visit:	20	21	22	23	24	25	26	27	Visit Type A ^{8, 10}	Visit Type B ⁹
Days on ATI:	D168	D196	D224	D252	D280	D308	D336	D364		
Weeks on ATI:	W24	W28	W32	W36	W40	W44	W48	W52		
Study procedures										
Screening consent (if used)										
Protocol consent										
Assessment of understanding										
Medical history										
Complete physical exam										
Targeted physical exam ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Concomitant medications ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercurrent illness/adverse experience ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
ART re-initiation assessment ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Contraception status assessment ⁵	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Decision aid										
Decision-making assessment	✓			✓				✓		✓
Psychosocial assessment	✓			✓				✓		✓
Social impact assessment ⁴	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Social impact assessment questionnaire	✓			✓				✓		✓
QuantiFERON tuberculosis test ⁶										
Confirm eligibility										
Specimen collection⁷	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ Screening may occur over the course of several contacts/visits up to and including the day of enrollment as defined in Section 6.1.2.

² For participants undergoing switch from NNRTI-based to protease- or integrase-based ART regimen. Participants undergoing an “ART switch” will be considered enrolled on first day of the new ART medication. The “ART switch” phase will not be performed for participants not on NNRTIs. For procedure timing during the “ART switch,” see Section 6.1.3 and HVTN 805/HPTN 093 SSP.

³ Enrollment visit for participants who do not undergo ART switch.

⁴ Procedure to be performed at interim visits held to draw confirmatory viral load samples or confirmatory samples for CD4+ T cell counts (see Sections 3.3.1, 3.3.2, and 3.3.3).

⁵ Contraception status assessment is required only for participants who can become pregnant (does not include those persons not of reproductive potential due to having undergone hysterectomy or bilateral oophorectomy [verified by medical records]).

⁶ If QuantiFERON TB testing cannot be performed (see Appendix E), TST should be conducted. Additional risk/clinical/diagnostic assessment may be performed at the discretion of the clinician to meet institutional standard of care for treatment of latent TB.

⁷ For specimen collection requirements, see Appendix E.

⁸ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 27, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁹ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 27, continuing up 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

¹⁰ At an early termination visit for a withdrawn or terminated participant, CRS staff should consider performing procedures specified for Extended follow-up visit Type A (see Section 6.5 and HVTN 805/HPTN 093 SSP for details).

Appendix I Procedures at CRS—Schedule 2: Monitoring ATI with viremia

Visit:	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55
Days on ATI with viremia:	D0	D7	D14	D21	D28	D35	D42	D49	D56	D70	D84	D98	D112	D126	D140	D154
Weeks on ATI with viremia:	W0	W1	W2	W3	W4	W5	W6	W7	W8	W10	W12	W14	W16	W18	W20	W22
Study procedures																
Complete physical exam	✓															
Targeted physical exam ¹		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Concomitant medications ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercurrent illness/adverse experience ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
ART re-initiation assessment ¹		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Contraception status assessment ²	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Decision-making assessment	✓															
Psychosocial assessment	✓										✓					
Social impact assessment ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Social impact assessment questionnaire											✓					
Specimen collection³	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ Procedure to be performed at interim visits held to draw confirmatory viral load samples or confirmatory samples for CD4+ T cell counts (see Sections 3.3.1, 3.3.2, and 3.3.3).

² Contraception status assessment is required only for participants who can become pregnant (does not include those persons not of reproductive potential due to having undergone hysterectomy or bilateral oophorectomy [verified by medical records]).

³ For specimen collection requirements, see [Appendix F](#).

⁴ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 66, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁵ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 66, continuing up 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

¹⁰ At an early termination visit for a withdrawn or terminated participant, CRS staff should consider performing procedures specified for Extended follow-up visit Type A (see Section 6.5 and HVTN 805/HPTN 093 SSP for details).

Procedures at CRS—Schedule 2: Monitoring ATI with viremia (continued)

Visit:	56	57	58	59	60	61	62	63	64	65	66	Visit Type A ^{4,6}	Visit Type B ⁵
Days on ATI with viremia:	D168	D182	D196	D210	D224	D238	D252	D280	D308	D336	D364		
Weeks on ATI with viremia:	W24	W26	W28	W30	W32	W34	W36	W40	W44	W48	W52		
Study procedures													
Complete physical exam													
Targeted physical exam ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Concomitant medications ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercurrent illness/adverse experience ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
ART re-initiation assessment ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Contraception status assessment ²	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Decision-making assessment	✓						✓				✓		✓
Psychosocial assessment	✓						✓				✓		✓
Social impact assessment ¹	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Social impact assessment questionnaire	✓						✓				✓		✓
Specimen collection³	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ Procedure to be performed at interim visits held to draw confirmatory viral load samples or confirmatory samples for CD4+ T cell counts (see Sections 3.3.1, 3.3.2, and 3.3.3).

² Contraception status assessment is required only for participants who can become pregnant (does not include those persons not of reproductive potential due to having undergone hysterectomy or bilateral oophorectomy [verified by medical records]).

³ For specimen collection requirements, see [Appendix F](#).

⁴ Extended follow-up visit type A will occur every 6 months starting with 3 months after visit 66, continuing up to 3 years of this schedule. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁵ Extended follow-up visit type B will occur every 6 months starting with 6 months after visit 66, continuing up 3 years of this schedule and then every 3 months thereafter. This follow-up visit may be performed for participants who have not met ART re-initiation criteria (see Protocol Section 3.3 and HVTN 805/HPTN 093 SSP for details).

⁶ At an early termination visit for a withdrawn or terminated participant, CRS staff should consider performing procedures specified for Extended follow-up visit Type A (see Section 6.5 and HVTN 805/HPTN 093 SSP for details).

Appendix J Procedures at CRS—Schedule 3: Follow-up on ART

Visit:	80 ¹	81	82	83	84	85	86	87	88	89	90	91 ⁴	92
Days after ART re-initiation:	D0	D14	D28	D42	D56	D70	D84	D112	D140	D168	D196	D280	D364
Weeks after ART re-initiation:	W0	W2	W4	W6	W8	W10	W12	W16	W20	W24	W28	W40	W52
Study procedures													
Complete physical exam													✓
Targeted physical exam	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Concomitant medications	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercurrent illness/adverse experience	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Transmission risk reduction counseling	✓	✓	✓		✓		✓		✓		✓		✓
Contraception status assessment ²	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Decision-making assessment	✓						✓			✓			✓
Psychosocial assessment	✓						✓			✓			✓
Social impact assessment	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Social impact assessment questionnaire							✓			✓			✓
Specimen collection³	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ ART re-initiation visit.

² Contraception status assessment is required only for participants who can become pregnant (does not include those persons not of reproductive potential due to having undergone hysterectomy or bilateral oophorectomy [verified by medical records]). Contraception status assessment is not required if participant VL has returned to undetectable.

³ For specimen collection requirements, see [Appendix G](#).

⁴ At an early termination visit for a withdrawn or terminated participant, CRS staff should consider performing procedures specified for Visit 92 (see Section [6.5](#) and HVTN 805/HPTN 093 SSP for more details).

Appendix K Visit Windows

Visit Windows – Schedule 1: Monitoring ATI

Visit Number	Visit Type		Lower Allowable Window	Lower Target Day	Target Day ⁴ <i>relative to initiation of ATI</i>	Upper Target Day	Upper Allowable Window
01.0 ¹	Screening [for participants <u>not</u> on NNRTIs]		-56	-	-	-	-
	Participants on NNRTIs [ART Switch Participants]	Screening ²	-140	-	-	-	-
		ART Switch ²	-84		-42		-29
02.0		ATI Qualification ^{2, 3}	-28		-14		0
03.0							
04.0	Week 0 Initiation of ATI		-	-	0	-	-
05.0	Week 1		-3	-2	7	+2	+3
06.0	Week 2		-3	-2	14	+2	+3
07.0	Week 3		-3	-2	21	+2	+3
08.0	Week 4		-3	-2	28	+2	+3
09.0	Week 5		-3	-2	35	+2	+3
10.0	Week 6		-3	-2	42	+2	+3
11.0	Week 7		-3	-2	49	+2	+3
12.0	Week 8		-3	-2	56	+2	+7
13.0	Week 10		-6	-4	70	+4	+7
14.0	Week 12		-6	-4	84	+4	+7
15.0	Week 14		-6	-4	98	+4	+7
16.0	Week 16		-6	-4	112	+4	+7
17.0	Week 18		-6	-4	126	+4	+7
18.0	Week 20		-6	-4	140	+4	+7
19.0	Week 22		-6	-4	154	+4	+7
20.0	Week 24		-6	-4	168	+4	+14
21.0	Week 28		-13	-8	196	+8	+14
22.0	Week 32		-13	-8	224	+8	+14
23.0	Week 36		-13	-8	252	+8	+14
24.0	Week 40		-13	-8	280	+8	+14
25.0	Week 44		-13	-8	308	+8	+14

26.0	Week 48	-13	-8	336	+8	+14
27.0	Week 52	-13	-8	364	+8	+14
Visit Type A ⁵	Follow-up (Schedule 1: Visit Type A)	-42	-28	Visit Type A will occur every 6 months starting with 3 months after visit 27 continuing up to 3 years of this schedule	+28	+42
Visit Type B ⁵	Follow-up (Schedule 1: Visit Type B)	-42	-28	Visit Type B will occur every 6 months starting with 6 months after visit 27 continuing up to 3 years of this schedule, and then every 3 months thereafter	+28	+42

¹Screening may occur over the course of several contacts/visits up to and including the day of enrollment, as defined in Section 6.1.2. All screening assessments must be done within 56 days of enrollment; for participants on NNRTIs, all screening assessments must be done within 56 days of the ART Switch.

²The ART switch phase will only be performed for participants on NNRTIs.

³Participants must be on the new ART regimen for at least 4 weeks prior to the ATI qualification visit, see Section 6.1.3. The ATI Qualification visit must be at least 28 days from the ART Switch and, if needed, VL interim visits may continue until approximately 12 weeks or until viral suppression has been achieved.

⁴Viral load sample collection should occur as close to the target day as possible and must occur at least 2 days apart (this applies to both scheduled visits and interim visits). See Visit Scheduling and Coding SSP for further details.

⁵Type A and Type B visits must be at least 8 weeks apart.

Visit Windows – Schedule 2: Monitoring ATI with viremia

Visit Number	Visit Type	Lower Allowable Window	Lower Target Day	Target Day ¹	Upper Target Day	Upper Allowable Window
40.0	First visit after confirmed ATI VL ≥ 200 copies/mL	-	-	0	-	-
41.0	Week 1	-3	-2	7	+2	+3
42.0	Week 2	-3	-2	14	+2	+3
43.0	Week 3	-3	-2	21	+2	+3
44.0	Week 4	-3	-2	28	+2	+3
45.0	Week 5	-3	-2	35	+2	+3
46.0	Week 6	-3	-2	42	+2	+3
47.0	Week 7	-3	-2	49	+2	+3
48.0	Week 8	-3	-2	56	+2	+7
49.0	Week 10	-6	-4	70	+4	+7
50.0	Week 12	-6	-4	84	+4	+7
51.0	Week 14	-6	-4	98	+4	+7
52.0	Week 16	-6	-4	112	+4	+7
53.0	Week 18	-6	-4	126	+4	+7
54.0	Week 20	-6	-4	140	+4	+7
55.0	Week 22	-6	-4	154	+4	+7
56.0	Week 24	-6	-4	168	+4	+7
57.0	Week 26	-6	-4	182	+4	+7
58.0	Week 28	-6	-4	196	+4	+7
59.0	Week 30	-6	-4	210	+4	+7
60.0	Week 32	-6	-4	224	+4	+7
61.0	Week 34	-6	-4	238	+4	+7
62.0	Week 36	-6	-4	252	+8	+14
63.0	Week 40	-13	-8	280	+8	+14
64.0	Week 44	-13	-8	308	+8	+14
65.0	Week 48	-13	-8	336	+8	+14
66.0	Week 52	-13	-8	364	+8	+14
Visit Type A ²	Follow-up (Schedule 2: Visit Type A)	-42	-28	Visit Type A will occur every 6 months starting with 3 months after visit 66 continuing up to	+28	+42

				3 years of this schedule		
Visit Type B ²	Follow-up (Schedule 2: Visit Type B)	-42	-28	Visit Type B will occur every 6 months starting with 6 months after visit 66 continuing up to 3 years of this schedule, and then every 3 months thereafter	+28	+42

¹Viral load sample collection should occur as close to the target day as possible and must occur at least 2 days apart (this applies to both scheduled visits and interim visits). See Visit Scheduling and Coding SSP for further details.

²Type A and Type B visits must be at least 8 weeks apart.

Visit Windows – Schedule 3: Follow-up on ART

Visit Number	Visit Type	Lower Allowable Window	Lower Target Day	Target Day¹ <i>relative to ART re-initiation</i>	Upper Target Day	Upper Allowable Window
80.0	ART Re-initiation	-	-	0	-	-
81.0	Week 2	-6	-4	14	+4	+7
82.0	Week 4	-6	-4	28	+4	+7
83.0	Week 6	-6	-4	42	+4	+7
84.0	Week 8	-6	-4	56	+4	+7
85.0	Week 10	-6	-4	70	+4	+7
86.0	Week 12	-6	-4	84	+4	+14
87.0	Week 16	-13	-8	112	+8	+14
88.0	Week 20	-13	-8	140	+8	+14
89.0	Week 24	-13	-8	168	+8	+14
90.0 ¹	Week 28	-13	-8	196	+8	+14
91.0 ¹	Week 40	-42	-14	280	+14	+42
92.0 ¹	Week 52	-28	-14	364	+14	+42

¹There must be at least 8 weeks between visits 90.0 and 91.0, and visits 91.0 and 92.0.

Appendix L Protocol Signature Page

Antiretroviral analytical treatment interruption (ATI) to assess immunologic and virologic responses in participants who initiated ART in early HIV infection after having received VRC01 or placebo in HVTN 703/HPTN 081

I will conduct the study in accordance with the provisions of this protocol and all applicable protocol-related documents. I agree to conduct this study in compliance with United States (U.S.) Health and Human Service regulations (45 CFR 46); applicable U.S. Food and Drug Administration regulations; standards of the International Conference on Harmonization Guideline for Good Clinical Practice (E6); Institutional Review Board/Ethics Committee determinations; all applicable in-country, state, and local laws and regulations; and other applicable requirements (eg, U.S. National Institutes of Health, Division of AIDS) and institutional policies

Investigator of Record Name (print)

Investigator of Record Signature

Date

DAIDS Protocol Number: HVTN 805/HPTN 093

DAIDS Protocol Version: Version 1.0

Protocol Date: April 27, 2020